

## 10.14. Surveys of Eagles and Other Raptors

### 10.14.1. General Description of the Proposed Study

The Surveys of Eagles and Other Raptors began in 2012 to prevent inadvertent take of raptors by providing information on raptor avoidance zones to Project personnel in the field in 2012. The Surveys of Eagles and Other Raptors will continue in 2013 and 2014, providing data for avoidance of raptor take, for the assessment of Project impacts, and ultimately for any necessary applications for federal eagle nest and take permits.

#### Study Goal and Objectives

The goal of the Surveys of Eagles and Other Raptors is to characterize population size, productivity, nesting phenology, and habitat use of raptor species to inform the prediction and quantification of impacts that may result from the proposed Project, and to provide information required for a possible application(s) for federal eagle take (lethal or disturbance take, see below) and/or eagle nest take permits. Common and scientific names of raptors that may occur in the Project area are listed in Table 10.14-1.

Six objectives have been identified for study:

- 1) Enumerate and identify the locations and status of raptor nests and territories that could be affected by Project construction and operations. Four specific tasks are associated with this objective:
  - a) Review and synthesize existing nest data for eagles and other raptors: Identify and assess the status of previously recorded nest locations of various species, including geographic coordinates, annual nest activity, descriptions of nest site characteristics, and general descriptions of cliff habitat in the proximity of each site.
  - b) Conduct field surveys to locate and characterize nests: Locate and map Bald Eagle and Golden Eagle nests in the Project study area, identifying all active and inactive nests and alternative nest sites. Locate and map active and inactive nests of other tree- and cliff-nesting raptor species (as well as Common Ravens) in the Project study area.
  - c) Create a geospatial database of all nests and territories: The database will be used to calculate inter-nest distances, estimate local average territory size, and, with overlays of Project footprint and habitats, determine the number of nests and territories potentially affected by the Project.
  - d) Calculate local average territory size for Bald Eagles and Golden Eagles: Estimates of average territory sizes (and mean inter-nest distance) are required for the applications for federal eagle nest take permits.
- 2) Estimate Project effects on productivity of raptors. This objective includes four specific tasks:
  - a) Review existing productivity data.
  - b) Determine the average and range of productivity of nests of each species (eagle/other raptor/raven).
  - c) Consider impacts on productivity at the local and larger population level using current and historical data.
  - d) Undertake pre- and post-construction comparisons of productivity to evaluate whether realized take is consistent with the permitted take, and to ensure that the level of take is compatible with the preservation of eagle populations.

- 3) Estimate effects on nesting and foraging habitats by delineating suitable habitat features in a geospatial database (this work will be conducted in the Evaluation of Wildlife Habitat Use; Section 10.19), and characterize and map habitat as suitable or not suitable for nesting and foraging for the various raptor species. These characterizations will be used in four ways:
  - a) Calculate the percentage of local habitat lost.
  - b) Calculate numbers of breeding pairs and productivity.
  - c) Estimate whether or not a partial loss of a territory may functionally result in abandonment of the entire territory.
  - d) Identify whether or not habitats adjacent to the Project area may be available for displaced nesting birds.
- 4) Conduct field surveys and literature reviews to identify, map, and characterize the habitat-use patterns at fall and winter communal roost sites and foraging sites of Bald and Golden eagles and other raptor species. Describe seasonal habitat use, highlighting areas or conditions that may result in impacts on raptors.
- 5) Conduct a study to assess the extent to which planned overhead transmission lines may pose a collision risk to migrating or nesting raptors and to identify migratory corridors (including altitudes of raptor movements) in the Project transmission line corridors.
- 6) Provide information on the distribution, abundance, food habits, and diet of piscivorous (fish-eating) raptors; feather samples for characterization of mercury levels; and information on the effects of methylmercury on piscivorous raptors, for use in the Mercury Assessment and Potential for Bioaccumulation study (see Section 5.7).

#### **10.14.2. Existing Information and Need for Additional Information**

Historical information from aerial surveys of raptors in the early 1980s provided the first assessment of the distribution, abundance, and vulnerability of many raptor nests located within the proposed Project impoundment zone. Those surveys highlighted Bald and Golden eagles and Common Ravens, and, to a lesser extent, raptors such as Northern Goshawks. Extensive information on raptors was collected during the 1980s for the original Alaska Power Authority (APA) Susitna Hydroelectric Project and for other surveys in the region (discussed in ABR 2011). Hard-copy maps are available of eagle nests located during the APA Susitna Hydroelectric Project studies in the early 1980s (LGL 1984). Other nest site records may exist in the files of the University of Alaska Museum of the North (AEA 2011). Other investigators and agency personnel may have information on raptor nest sites and important habitats, such as roosting sites, in or near the Project area. Similar regional databases of nest site information have been developed (Wildman and Ritchie 2000).

Surveys completed in the middle and upper Susitna River valley during the 1980s identified 23 Golden Eagle, 10 Bald Eagle, 3 Gyrfalcon, 3 Northern Goshawk, and 21 Common Raven nest sites (some sites included more than one nest site, if they were close together) (APA 1985). Although Common Ravens are not raptors, they construct both cliff and tree nests that raptors often use, are culturally significant, and are protected by the Migratory Bird Treaty Act (MBTA). Of the eagle nest sites identified in the 1980s, 5 Golden Eagle and 3 Bald Eagle sites were expected to be completely inundated by the original APA Susitna Hydroelectric Project Phase I (Low Watana) impoundment (LGL 1984).

New raptor studies are needed because most of the existing information is almost 30 years old and it is unknown how distribution, status, or other conditions may have changed. Also,

historical surveys did not cover the entire area of current interest, including access roads and power transmission corridors. More sophisticated geospatial analyses are now available that allow for more accurate assessments of the potential effects of the Project on raptors and their habitats. Finally, current data will be necessary for compliance with federal laws, especially the Bald and Golden Eagle Protection Act (BGEPA) and the MBTA, the 2011 FERC–USFWS Memorandum of Understanding (MOU), and application(s) for federal eagle nest/territory take permits.

A limited field survey for raptors was conducted in 2011 (ABR 2011) and more extensive surveys of the Project area were conducted in 2012 (AEA 2012) to provide current information needed to protect raptors by restricting Project activities near active raptor nests during pre-license field studies and construction. In 2011, surveys on June 27 were limited to the area near borehole sites drilled for the geotechnical program in the vicinity of the proposed Watana Dam. In 2012, occupancy surveys for nesting raptors were performed twice in May and productivity surveys were performed twice in July. The 2012 survey area comprised the area within a 2-mile buffer of the Project area (impoundment, access and transmission corridors, and facilities/infrastructure). Surveys were conducted from a Robinson R44 piston helicopter. Dozens of raptor nests were observed and occupied nest sites were located and mapped. In addition to Common Ravens, nests of six species of raptors were identified in the Project area in 2012: Golden Eagle, Bald Eagle, Peregrine Falcon, Gyrfalcon, Red-tailed Hawk, and Merlin. Global Positioning System (GPS) files, maps, and avoidance guidelines were distributed to Project personnel and contractors to avoid “take” of nests by disturbance.

Although transmission lines can be a source of mortality for eagles and other raptors by electrocution and collision, it is assumed that all new transmission lines and power transfer stations for the Project will be built to the “eagle-safe” standards developed by the Avian Power Line Interaction Committee (APLIC 2006), and therefore will not be likely to constitute a significant source of electrocution risk for raptors. However, significant lengths of new transmission lines will be constructed across miles of open and undisturbed landscape. As discussed in the Avian Protection Plan (APP) Guidelines (APLIC and USFWS 2005), collision risk assessments are recommended in the siting of overhead power transmission lines.

### **10.14.3. Study Area**

The study area is subdivided into two different-sized survey areas, depending on the species of interest. For Golden Eagles, 10 miles is the survey radius typically recommended by the U.S. Fish and Wildlife Service (USFWS) for Golden Eagles (Pagel et al. 2010) in areas that contain suitable nesting habitat. After consultation with AEA, however, USFWS agreed to a 10-mile survey radius only around the reservoir inundation zone to calculate a mean inter-nest distance for this species and a 3-mile radius around proposed facilities and potential access road and transmission corridors (M. de Zeeuw and J. Muir, USFWS, pers. comm., April 2012).

For Bald Eagles and other raptor species, a radius of 3 miles around the reservoir inundation zone, proposed facilities, and centerlines of the potential access road and transmission line corridors is considered to be sufficient (M. de Zeeuw and J. Muir, USFWS, pers. comm., April 12, 2012).

Hence, the study area for the Survey of Eagles and Other Raptors encompasses: (1) a 10-mile radius around the reservoir inundation zone for Golden Eagles; (2) a 3-mile radius around the

reservoir inundation zone for Bald Eagles and other raptor species; and (3) a 3-mile buffer for all eagles and other raptors around proposed facilities and the centerlines of the potential access road and transmission line corridors (Figure 10.14-1).

All Bald and Golden eagle habitat within the relevant survey area boundaries will be surveyed. For Bald Eagles, surveys will cover the area within a half-mile of the centers of all drainages with suitable timber and within a half-mile of all shorelines of lakes with similar characteristics in the inundation zone and wherever these habitats cross proposed road and transmission line corridors. Information on other large tree-nesting birds will also be collected during those surveys. Survey routes for cliff-nesting raptors will be flown in a cliff-to-cliff survey pattern, focusing on cliffs suitable for Golden Eagle nests.

The survey methodology will obtain information for an area larger than the 1980s survey coverage, will gather information on key species in a more well-defined study impact area, and will provide information needed for eagle permitting and to develop avoidance areas and mitigation protocols to reduce the potential disturbance of nesting raptors from Project construction and operations. The nesting survey may be sectioned to include segments that match the extent of the 1980s survey to the extent appropriate for comparative purposes to evaluate trends in raptor populations and/or habitat use.

The study area for migration route surveys may be limited to specific locations along planned transmission line routes that may pose risks to migrating birds (e.g., ridgelines). These study areas will be determined in consultation with USFWS and be based on review of existing raptor migration data, topographical and wind current information, and other relevant factors.

#### **10.14.4. Study Methods**

##### *10.14.4.1. Field Surveys*

Inventory and monitoring methodologies for nest occupancy and productivity surveys will follow established aerial and ground-based protocols for eagle nest surveys (USFWS 2007; Pagel et al. 2010), using appropriately trained observers and suitable survey platforms (helicopter or fixed-wing aircraft). Modifications may be necessary to extend to the objective of identifying and monitoring the nests of other raptors. Nests of cliff-nesting raptors (including Golden Eagle, Peregrine Falcon, Gyrfalcon, and potentially Bald Eagle) and raptors using large stick nests (including Bald Eagle, Great Horned Owl, Northern Goshawk, Red-Tailed Hawk, Osprey, and potentially Golden Eagle) will be inventoried and monitored, as will raven nests.

Small to medium-sized raptor species (e.g., Short-eared Owl, Boreal Owl, Northern Hawk Owl, Northern Harrier, American Kestrel, Merlin, and Sharp-shinned Hawk) will be recorded during ground-based surveys for the breeding landbird and shorebird point-count surveys (see Section 10.16).

Raptor nest occupancy surveys will begin in spring before leaf-out (late April to late May), focusing on primary habitats for Bald and Golden eagles, but also considering primary habitats of resident species nesting in woodlands (e.g., Great Horned Owl and Northern Goshawk) and on cliffs (e.g., Gyrfalcon and Peregrine Falcon).

The nest productivity survey period will extend from mid-June to late July to verify and monitor nesting activity and to search for additional nests of later-nesting raptors. Because of the wide

range of breeding dates for all raptors considered in the study (mid-February for resident owls through early September for dispersal of Bald Eagles from nesting areas), the second survey period will encompass a broad timing window. The nesting chronology of each focal species of raptor will be considered during survey scheduling.

The same helicopter protocols will be employed for the occupancy and productivity surveys. A helicopter will be used, carrying two observers in addition to the pilot. Flight altitude and speed will follow standard survey protocols for each habitat type (Pagel et al. 2010). Observers will be seated on the same side of the aircraft during surveys. Location and nest attribute data, including substrate, species, and nest status, will be collected for entry into a geodatabase.

In any aerial survey, a key concern is quantifying the sightability of the target species to adjust density estimates for targets missed. The actual sightability of nests depends on many factors, including nest size, location, survey weather/light conditions, substrate and tree density, habitat type, observer experience, and survey platform. Although Golden and Bald eagles often construct large, conspicuous stick nests, some inconspicuous nests are likely to be missed when conducting surveys. Re-surveys of sub-samples of the study area will be performed to quantify the sightability of raptor nests in the Project area.

To prevent disturbance to Dall's sheep during the lambing period and near the Jay Creek and Watana Creek mineral lick sites, standard eagle survey protocols may need to be modified (Pagel and Whittington 2011) and helicopter surveys will be routed to avoid these areas during these periods.

The wildlife habitat map developed by the Vegetation and Wildlife Habitat Mapping Study in the Upper and Middle Susitna Basin (Section 11.5) will be used to delineate the probable distribution of early nesting owls. Results from point counts conducted for the Landbird and Shorebird Migration, Breeding, and Habitat Use study (Section 10.16) will be combined with information from a literature review to assess the distribution, abundance, and habitat use by these owl species, which will then be incorporated into the Evaluation of Wildlife Habitat Use (Section 10.19). No winter surveys are proposed for early nesting owls because they would require logistically difficult and potentially dangerous nocturnal surveys during winter in remote areas to obtain information on only two species (Boreal Owl and Northern Hawk Owl) that are uncommon. Great Horned and Great Gray owl nests will be recorded during aerial surveys for Bald Eagles and other tree-nesting raptors.

Surveys for foraging and communal roost locations will be conducted primarily in fall and early winter. Repeated surveys of suitable protected forest stands may be necessary due to the high mobility of wintering Bald Eagles. Four aerial surveys of foraging habitat and communal roosts, primarily for Bald Eagles, will be conducted each year at intervals of 7 to 21 days between mid-October and early December. Survey numbers and timing may be adjusted in 2013 and 2014, based on the results of the initial surveys conducted in 2012. A helicopter or a fixed-wing aircraft carrying two observers will be used for these surveys. Surveys will be conducted near dawn or dusk. Information on fall fishery concentrations will be requested from Project fisheries researchers and from agency biologists to more effectively monitor potential Bald Eagle concentration areas.

Surveys to assess whether migrating raptors would be at risk for collision with Project power transmission lines will be conducted using fixed-radius migration point counts. These surveys generally will follow the USFWS's recommended point-count protocol, based on the standard

hawk migration counting protocols described in Appendix C of the *Draft Eagle Conservation Plan Guidance* (USFWS 2011). Migration point counts will be centered in plots with a radius of 800 meters and will be conducted for a period of 30 minutes each. The plot locations will be delineated along the transmission corridor alternatives before surveys begin, and the final selection of plots to be sampled will be determined by focusing on areas judged likely to concentrate migrating raptors (on the basis of topography). Survey efforts will be timed to coincide with times of day when thermal updrafts are most likely to occur (from midday through the afternoon hours).

After federal and state permits have been acquired and nests of eagles and other raptors have been vacated for the season, a sample of nests of piscivorous raptors (primarily Bald Eagle but also Osprey, if any nests of the latter species are found) in the vicinity of the proposed reservoir will be visited to obtain samples of feathers for laboratory analysis of mercury levels, which will be provided to the investigators conducting the Mercury Assessment and Potential for Bioaccumulation Study (Section 5.7). A literature review will be conducted for this study to provide basic information on the food habits and diets of, as well as the documented effects of methylmercury on, piscivorous raptors. Because this information will be provided to the investigators conducting the mercury study, it will not be included in the reports prepared for the study of eagles and other raptors.

#### 10.14.4.2. Reporting

Reporting of inventory and monitoring data will comply with the protocols and standards described in the MOU between FERC and USFWS (FERC and USFWS 2011). Survey reports will include the following:

- Maps and associated metadata for historical eagle and other raptor nest and communal roost locations with survey extents to compare to current survey data.
- Maps and associated metadata with coordinates for current nest locations, nest activity status, fall and winter communal roost areas, and migration routes.
- Summary and mapping of suitable forest, riparian, and cliff habitats to evaluate the extent of suitable nesting habitats and facilitate nest searches within the study area.

Observations will be recorded and geo-referenced with associated habitats during raptor surveys. Some raptor nests and observations will also be recorded during breeding landbird and shorebird point-count surveys. All raptor observations will be plotted on the wildlife habitat map using field GPS coordinates. Nest characteristics will be recorded according to a protocol developed in consultation with the USFWS, including the protocol of the USFWS Alaska Bald Eagle Nest Atlas (<http://alaska.fws.gov/mbsp/mbm/landbirds/alaskabaldeagles/default.htm>).

The wildlife habitat map will provide the basis for an ecosystem approach to assessing the effects of development-related habitat impacts on raptors. The wildlife habitat map will facilitate quantitative spatial analyses of raptor habitat availability and changes likely to result from development, and, in combination with raptor survey data, will provide a way to assess the potential for changes in local raptor populations during construction and operations. Spatial analyses will be used to calculate the area and percentage of habitat lost and the numbers of breeding pairs and their productivity that likely would be affected; to determine whether or not a partial loss of a given territory may functionally result in abandonment or failure of the entire territory; to identify whether or not habitats adjacent to the Project area may be “available”

(notwithstanding occupancy) for displaced nesting birds; and to assess the risk of raptor collisions with overhead transmission lines.

#### 10.14.4.3. Data Analysis

A geospatially referenced relational database will be developed to incorporate historical and current data, including nest and roost locations for each species; occupancy, activity, and productivity data; nest type and characteristics; vegetation stand characteristics; and photographs. Suitable raptor nesting habitat will be delineated using a combination of field mapping and Geographic Information System (GIS) analysis. Existing nest locations and distribution of timber stands with suitably sized nest trees, in coordination with Project studies involving vegetation surveys and mapping, will be incorporated into the identification and delineation of suitable raptor nesting habitats. Foraging habitats will also be delineated whenever possible. Distribution of spawning salmon (determined through collaboration with Project salmon studies, Sections 9.5, 9.6, and 9.7) will be used to identify Bald Eagle foraging locations and potential aggregation areas. Distribution of fall waterfowl staging areas (determined in coordination with the waterbirds study, Section 10.15) will provide additional information for locating fall Bald Eagle foraging locations and potential communal roost areas. The distribution of Dall's sheep lambing areas and caribou calving areas, identified in part by the studies of Dall's Sheep Distribution and Abundance (Section 10.7) and Caribou Distribution, Abundance, Movements, Productivity, and Survival (Section 10.6), will provide information for Golden Eagle foraging habitat analyses.

Local Bald Eagle and Golden Eagle territory sizes will be estimated using inter-nest distances, as described in the *Draft Eagle Conservation Plan Guidance* (USFWS 2011). Recommendations will be developed for future data-gathering needs and analyses designed to evaluate potential Project-related impacts to eagles and other raptors.

#### 10.14.4.4. Deliverables

Study products will include the following:

Geospatially-Referenced Relational Database. A geospatially-referenced relational database will be developed that incorporates all historic and current data, including nest, forage, and roost locations for each species; occupancy/activity; nest type and characteristics; stand characteristics; and photographs. This database will be expanded from the work done for the 2012 Raptor Study. All field data must be associated with location information collected using a GPS receiver in un-projected geographic coordinates (latitude/longitude) and the NAD 83 datum (or convertible as such). Migratory corridor information will be included for specific areas of concern, as discussed above.

Delineation of Suitable Eagle and Raptor Nesting and Foraging Habitats. Habitat delineation will be completed using GIS software as part of the Vegetation and Wildlife Habitat Mapping Study (Section 11.5) and the Evaluation of Wildlife Habitat Use (Section 10.19).

Study Reports. The Initial Study Report will be completed February 2014 and the Updated Study Report will be completed February 2015. The Updated Study Report will summarize the results for both years (plus 2012). These reports will include the following information:

- Discussion of nest-mapping results

- Calculation of the following:
  - Local average territory size for Bald Eagle and Golden Eagle
  - Productivity (annual, mean, range) for each raptor species and Common Raven among Project subareas (reservoir impoundment zone, access roads, power transmission corridors)
- Discussion of migration survey results
- Preliminary discussions and calculations of potential Project impacts, including the following:
  - Numbers of nests and territories that will be lost per species per subarea
  - Numbers of nests and territories otherwise affected per subarea
  - Type and level of impacts to forage and roost areas
  - Locations of any potential collision hazard areas for migrating raptors
  - Other potential impacts, including large increases in open-water habitats created by the reservoir impoundment

#### **10.14.5. Consistency with Generally Accepted Scientific Practice**

The study methods described above are consistent with generally accepted scientific practice. The field protocols may be modified to address logistical constraints imposed by the size and remoteness of the study area. The field protocols for raptor surveys generally follow established techniques for cliff- and tree-nesting raptors in North America (e.g., Anderson 2007). In addition, survey protocols and study areas will be tailored for specific species. For example, inventory and monitoring methodologies for nest occupancy and productivity surveys follow established aerial and ground-based protocols for eagle nest surveys (USFWS 2007; Pagel et al. 2010), using appropriately trained observers and suitable survey platforms (helicopter or fixed-wing aircraft). Nest characteristics will be recorded according to protocols developed in consultation with USFWS, including the protocol of the USFWS Alaska Bald Eagle Nest Atlas (<http://alaska.fws.gov/mbmp/mbm/landbirds/alaskabaldeagles/default.htm>). Local Bald Eagle and Golden Eagle territory sizes will be estimated using inter-nest distances as described in the *Draft Eagle Conservation Plan Guidance* (USFWS 2011). Surveys to determine if migration routes exist that may put migrating raptors at risk for collision with Project power transmission lines will generally follow USFWS's recommended migration point-count protocol, based on standard hawk migration counting protocols described in Appendix C of the *Draft Eagle Conservation Plan Guidance* (USFWS 2011).

#### **10.14.6. Schedule**

This study is a multi-year effort that began in 2012 and will continue through 2013 and 2014 into the first quarter of 2015. The data-gathering and reporting schedule is described in more detail in Table 10.14-2.

Nest occupancy and productivity surveys will be conducted periodically between late April and late July in 2013 and 2014. Occupancy surveys will be conducted between late April and late May and productivity surveys will be conducted between mid-June and late July. A minimum of two aerial surveys at least 30 days apart are recommended by USFWS for the Golden Eagle protocol (Pagel et al. 2010). Early reporting of potentially active raptor nest sites after the initial surveys in May (potentially earlier, depending on USFWS recommendations) will be used to develop avoidance measures for Project-related field study activities that have the potential to



disturb active nests. As soon as they are found, active eagle and other raptor nest sites will be reported to AEA in order to develop avoidance zones for field studies.

Raptor migration point-count surveys will be conducted during April–May and September–October in 2013 and 2014. Surveys will be conducted during peak periods of raptor migration in spring and fall.

Field survey data will be used to update the geospatially referenced, relational database of historical and current nest data in August 2013 and 2014, after occupancy and productivity surveys have been completed. At that time, the most current delineation of suitable eagle and raptor nesting habitat and the locations of active and inactive nest locations will be entered into the database and proofed.

Roosting and staging surveys will be conducted between mid-October and early December in 2013 and 2014. Surveys will be conducted periodically to identify the use of winter foraging and potential communal roost sites along the Susitna River. Four aerial surveys will be flown at intervals of 2 to 3 weeks, depending on weather and the results of preceding surveys.

The Initial Study Report and Updated Study Reports will be completed within 1 and 2 years, respectively, of FERC's Study Plan Determination (February 1, 2013). The study reports will include a summary of the study results to date. In addition, study updates will be presented at the Technical Workgroup meetings, which will be held quarterly during 2013 and 2014.

#### **10.14.7. Relationship with Other Studies**

Information from two other studies will provide useful input for planning surveys of eagles and other raptors (Figure 10.14-2). Preliminary mapping of forested habitats from the Vegetation and Wildlife Habitat Mapping Study in the Upper and Middle Susitna Basin (Section 11.5) will help in survey planning for Bald Eagles and other tree-nesting raptors. Information on the location and timing of late-season spawning runs of salmon from the studies of Fish Distribution and Abundance in the Upper Susitna River (Section 9.5), Fish Distribution and Abundance in the Middle and Lower Susitna River (Section 9.6), and the Salmon Escapement Study (Section 9.7) will help in planning surveys of roosting and staging eagles in the fall and early winter. The geospatial database of raptor nest locations will contribute directly to the Vegetation and Wildlife Habitat Mapping Study in the Upper and Middle Susitna Basin (Section 11.5) and to the Bat Distribution and Habitat Use study (Section 10.13) by identifying suitable cliff-nesting habitats. Seasonal location and distribution data from the nest occupancy and productivity surveys, roosting and staging surveys, and migration surveys will be used to identify high-value wildlife habitat types for different raptor species in the Evaluation of Wildlife Habitat Use (Section 10.19). Data on nest distribution, species abundance, and productivity, as well as data on the numbers and flight paths of raptors during migration surveys of the transmission corridors, will be used to inform the assessment of Project impacts in 2015 in the FERC License Application and to help identify any potential PM&E measures, as appropriate. Feathers collected from nests after the breeding season will provide samples to characterize pre-construction levels of mercury in tissues of piscivorous raptors for the Mercury Assessment and Potential for Bioaccumulation study (Section 5.7).

The primary impact mechanisms of the Project on raptors may include the following:

- Permanent direct and indirect habitat loss and alteration, including loss of nesting sites and loss and alteration of foraging habitat.
- Temporary direct and indirect habitat loss and alteration, including indirect impacts resulting from altered distribution and abundance of prey.
- Potential direct behavioral impacts, such as attraction or avoidance, resulting from vehicular use, noise, and increased human presence associated with Project construction or operation.
- Potential indirect behavioral impacts to wildlife, such as attraction or avoidance, resulting from changes in vehicular use, noise, and increased human presence associated with increased subsistence or recreational access that may be facilitated by Project development.
- Potential direct mortality due to strikes with vehicles, power lines, towers, or other Project facilities; exposure to contaminants; and attraction to garbage and human activity.

Any impacts associated with habitat loss and alteration, attraction and avoidance, and direct mortality will occur primarily in the Project area, including the reservoir inundation zone, access and transmission corridors, and other Project infrastructure. Impacts associated with altered distribution and abundance of prey may occur over a larger area due to potential changes in both competing mammalian predators and prey species abundance.

During the impact assessment that will be conducted in 2015 for the FERC License Application, the impacts of direct and indirect habitat loss and alteration can be assessed through geospatial analysis. When plotted on the wildlife habitat map, raptor nest location data will allow the identification of high-value breeding habitats. Similarly, important habitats for prey species will also be identified by the Evaluation of Wildlife Habitat Use (Section 10.19). Using GIS software, the direct impacts of habitat loss can be evaluated for each raptor species by overlaying the reservoir impoundment, related infrastructure areas, and access road and power transmission corridors onto the wildlife habitat map prepared by the Vegetation and Wildlife Habitat Mapping in the Upper and Middle Susitna Basin study (Section 11.5) to calculate loss of preferred habitats. As was noted earlier, pertinent data from other studies will be incorporated into the evaluation of potential Project-related impacts to eagles and other raptors. Indirect impacts of habitat loss and alteration and behavioral reactions (such as avoidance) can be estimated by applying various buffer distances, as determined from the literature on the effects of similar projects, including responses of both raptor and their prey. In this way, the GIS analysis can be combined with information from the literature to estimate the geographic extent, frequency, duration, and magnitude of Project effects on raptor populations. Effects on the habitats of prey species included in the Evaluation of Wildlife Habitat Use (Section 10.19) can be incorporated into the impact assessment for raptors. Any necessary protection, mitigation, and enhancement (PM&E) measures will be developed, as appropriate by examining the distribution and abundance of raptor species and habitats in relation to the geographic extent and seasonal timing of various Project activities.

Data collected for this study will allow calculation of the numbers of nests and territories that will be lost per species per sub-area; the numbers of nests and territories otherwise affected per sub-area; the type and level of impacts to forage and roost areas; the locations of any potential collision hazard areas for migrating raptors; and other potential impacts, including large increases in the availability of open water habitats created by the impoundment.

### 10.14.8. Level of Effort and Cost

Occupancy/productivity for nesting raptors and fall and winter roost/forage surveys in 2013–2014 will require an estimated 10–12 days of additional fieldwork beyond the 2012 surveys due to the extended survey area for Golden Eagles. Therefore, costs for these surveys (including helicopter time, analysis, and reporting) will be approximately \$500,000 per year.

Transmission corridor surveys for migrating raptors in 2013–2014 will require approximately 30 field days, and estimated costs for these (with helicopter drop-offs, literature search, analysis, and reporting) will be approximately \$80,000 per year.

### 10.14.9. Literature Cited

- ABR. 2011. Wildlife data-gap analysis for the proposed Susitna-Watana Hydroelectric Project. Draft report, August 16, 2011. Report for the Alaska Energy Authority by ABR, Inc.—Environmental Research & Services, Fairbanks, Alaska. 114 pp.
- ADF&G (Alaska Department of Fish and Game). 2006. *Our wealth maintained: A strategy for conserving Alaska's diverse wildlife and fish resources*. Alaska Department of Fish and Game, Juneau, Alaska. 842 pp.
- AEA (Alaska Energy Authority). 2011. Pre-application document: Susitna-Watana Hydroelectric Project, FERC Project No. 14241. December 29, 2011. Prepared for the Federal Energy Regulatory Commission, Washington, D.C.
- AEA. 2012. Surveys of eagles and other raptors for the Susitna–Watana Hydroelectric Project, FERC Project No. 14241. Draft final version (March 26, 2012). Alaska Energy Authority, Anchorage.
- Anderson, D. E. 2007. Survey techniques. Pp 89-100 *in: Raptor Research and Management Techniques* (D. Bird and K. Bildstein, eds.). Hancock House Publishers LTD, Surrey, B.C. Canada. 463 pp.
- APA (Alaska Power Authority). 1985. Before the Federal Energy Regulatory Commission: Draft amended application for license for major project—Susitna Hydroelectric Project. Volume 11, Exhibit E – Chapter 3: Fish, wildlife, and botanical resources. 702 pp. [APA Doc. No. 3435]
- APLIC (Avian Power Line Interaction Committee). 2006. *Suggested practices for avian protection on power lines: The state of the art in 2006*. Edison Electric Institute, APLIC, and the California Energy Commission, Washington, D.C., and Sacramento, CA.
- APLIC and USFWS. 2005. Avian Protection Plan (APP) Guidelines. <http://www.aplic.org/mission.php>
- BLM (Bureau of Land Management). 2010. BLM–Alaska sensitive animal and plant lists. Alaska State Office, Anchorage.
- BPIFWG (Boreal Partners in Flight Working Group). 1999. Landbird conservation plan for Alaska biogeographic regions. Version 1.0. U.S. Fish and Wildlife Service, Anchorage.
- FERC and USFWS. 2011. Memorandum of Understanding between the Federal Energy Regulatory Commission and the U.S. Department of the Interior United States Fish and

- Wildlife Service Regarding Implementation of Executive Order 13186, “Responsibilities of Federal Agencies to protect Migratory Birds.” March 2011. 13 pp.
- Kessel, B., S. O. MacDonald, D. D. Gibson, B. A. Cooper, and B. A. Anderson. 1982. Susitna Hydroelectric Project environmental studies, Phase I final report – Subtask 7.11: Birds and non-game mammals. Report prepared for the Alaska Power Authority by the University of Alaska Museum, Fairbanks, and Terrestrial Environmental Specialists, Inc., Phoenix, New York. 149 pp. [APA Doc. Nol. 398.]
- LGL. 1984. Susitna Hydroelectric Project: Update and refinement of Bald and Golden eagle impact assessments and mitigation plans. Report for the Alaska Power Authority by LGL Alaska Research Associates, Inc., Anchorage. 68 pp. [APA Doc. No. 2374]
- Pagel, J. E., D. M. Whittington, and G. T. Allen. 2010. *Interim Golden Eagle technical guidance: Inventory and monitoring protocols and other recommendations in support of eagle management and permit issuance*. U.S. Fish and Wildlife Service, Division of Migratory Bird Management, Washington, DC. 30 pp.
- Pagel, J. E., and D. M. Whittington. 2011. *Interim Golden Eagle Inventory and Monitoring Protocols; and Other Recommendations: 2011*. Presented at the Western Raptor Symposium. J. L. Lincer and D. Bittner, co-chairs. Hosted by Wildlife Research Institute and The Wildlife Society, Western Section. Riverside Convention Center, Riverside, California. February 8–9, 2011.
- USFWS (U.S. Fish and Wildlife Service). 2007. *National Bald Eagle management guidelines*. Washington, DC. 25 pp.
- USFWS. 2008. *Birds of Conservation Concern, 2008*. Division of Migratory Bird Management, Arlington, Virginia. 85 pp. Available online (accessed November 2011) at: <http://www.fws.gov/migratorybirds/NewReportsPublications/SpecialTopics/BCC2008/BCC2008.pdf>
- USFWS. 2009. 50 CFR Parts 13 and 22. Eagle permits; take necessary to protect interests in particular localities; final rules. *Federal Register* 74: 46,836–46,879.
- USFWS. 2011. *Draft Eagle Conservation Plan Guidance*. Available online at: [http://www.fws.gov/windenergy/eagle\\_guidance.html](http://www.fws.gov/windenergy/eagle_guidance.html).
- Wildman, A., and R. J. Ritchie. 2000. Synthesis of survey information on cliff-nesting raptors and their habitats on the North Slope with emphasis on Peregrine Falcons in the Eastern NPR-A and recommendations for survey needs. Report to U.S. Fish and Wildlife Service, Northern Alaska Ecological Services, Fairbanks, by ABR, Inc.—Environmental Research & Services, Fairbanks.

### 10.14.10. Tables

**Table 10.14-1. Raptors in the vicinity of the middle basin of the Susitna River (from Tables 4.6-2 and 4.8-2 in AEA 2011).**

Common Name	Scientific Name	Conservation Status <sup>1</sup>	Seasonal Status <sup>2</sup>	Relative Abundance <sup>3</sup>
Bald Eagle	<i>Haliaeetus leucocephalus</i>	FS	B	uncommon
Boreal Owl	<i>Aegolius funereus</i>	PIF, FS	R	rare
Golden Eagle	<i>Aquila chrysaetos</i>	BLM, FS	B	fairly common
Great Gray Owl	<i>Strix nebulosa</i>	PIF, FS	?	rare
Great-horned Owl	<i>Bubo virginianus</i>	FS	R	uncommon
Gyrfalcon	<i>Falco rusticolus</i>	PIF, FS	R	uncommon
Merlin	<i>Falco columbarius</i>	FS	B	uncommon
Northern Harrier	<i>Circus cyaneus</i>	FS	B	fairly common
Northern Goshawk	<i>Accipiter gentilis</i>	FS	B	uncommon
Northern Hawk Owl	<i>Surnia ulula</i>	FS	R	uncommon
Osprey	<i>Pandion haliaetus</i>	FS	M	rare
Peregrine Falcon	<i>Falco peregrinus anatum</i>	BCC, FS	M	unknown
Red-tailed Hawk	<i>Buteo jamaicensis</i>	FS	B	uncommon
Short-eared Owl	<i>Asio flammeus</i>	BLM, FS	B?, M, S	uncommon
Sharp-shinned Hawk	<i>Accipiter striatus</i>	FS	B	uncommon

Notes:

- 1 Conservation Status: FS = Featured Species (ADF&G 2006); BCC = Birds of Conservation Concern (USFWS 2008); BLM = BLM Sensitive Species (BLM 2010); PIF = Boreal Partners in Flight Working Group (BPIFWG 1999).
- 2 Seasonal Status: M = migrant (transient); B = breeding; S = summering; R = resident; ? = uncertain (Kessel et al. 1982; APA 1985: Appendices E5.3 and E6.3).
- 3 From Kessel et al. (1982) and APA (1985: Appendices E5.3 and E6.3).

**Table 10.14-2. Schedule for implementation of Surveys of Eagles and Other Raptors.**

Activity	2012			2013			2014				2015		
	2 Q	3 Q	4 Q	1 Q	2 Q	3 Q	4 Q	1 Q	2 Q	3 Q	4 Q	1 Q	
Aerial surveys for nest occupancy and productivity assessments (2 surveys each)	—				—				—				
Migration surveys (transmission corridors)					—		—						
Update geospatial database of historical and current nesting data							—				—		
Update delineation of suitable nesting habitat, old and active nest locations, historical fall and winter roost locations					—	—	—		—	—	—		
Conduct roosting and staging surveys			—				—				—		
Initial Study Report								—	△				
Updated Study Report												—	▲

Legend:

- Planned Activity
- △ Initial Study Report
- ▲ Updated Study Report

10.14.11. Figures




Figure 10.14-1. Study area for Surveys of Eagles and Other Raptors.

**STUDY INTERDEPENDENCIES FOR SURVEYS OF EAGLES & OTHER RAPTORS**




Figure 10.14-2. Study interdependencies for Surveys of Eagles and Other Raptors.



## **10.15. Waterbird Migration, Breeding, and Habitat Use Study**

### **10.15.1. General Description of the Proposed Study**

The Waterbird Migration, Breeding, and Habitat Use Study will be conducted over two years (2013 and 2014) and will include aerial surveys of water bodies during spring and fall migration, a study of diurnal and nocturnal migration using visual and radar sampling, breeding-pair surveys, stream surveys for Harlequin Ducks, and brood-rearing surveys. Waterbirds may use lakes, ponds, rivers, and flooded wetland areas throughout the Project area to varying degrees during spring and fall migration. Aerial surveys for staging and migration will follow a lake-to-lake pattern and will also parallel river courses. The migration study will employ intensive monitoring of migrating birds during both daytime and nighttime hours at a site located near the proposed dam and associated camp infrastructure. Surveys of breeding waterbirds (primarily waterfowl) will use a combination of full-coverage lake-to-lake surveys in most of the study area and breeding-pair transect surveys in the easternmost portion of the study area. Aerial surveys for Harlequin Ducks will focus on river and stream habitats during the pre-nesting and brood-rearing seasons. Brood-rearing surveys will be conducted by surveying open water and shoreline habitats of lakes and ponds in the study area.

#### **Study Goals and Objectives**

The goal of the Waterbird Migration, Breeding, and Habitat Use Study is to collect baseline data on waterbirds migrating through and breeding in the Project area to enable assessment of the potential impacts of the Project and to inform the development of appropriate protection, mitigation, and enhancement measures. As used here, “waterbirds” is applied broadly to include swans, geese, ducks, loons, grebes, cranes, cormorants, herons, gulls, and terns. Shorebirds frequently are included in the general category of waterbirds, but they are addressed separately for this Project under the Landbird and Shorebird Migration, Breeding, and Habitat Use Study (Section 10.16) because the ground-based survey methods for shorebirds are similar to those used for landbirds. This study plan includes breeding surveys for the Harlequin Duck, a species of conservation concern that requires specific stream-survey techniques.

This study has three objectives:

- Document the occurrence, distribution, abundance, habitat use, and seasonal timing of waterbirds migrating through the Project area in spring and fall.
- Document the occurrence, distribution, abundance, productivity, and habitat use of waterbirds breeding in the Project area.
- Review available information to characterize food habits and diets of piscivorous waterbirds documented in the study area as background for the Mercury Assessment and Potential for Bioaccumulation Study (Section 5.7).

The information gained from this study will be used to evaluate waterbird habitat loss and alteration quantitatively, in conjunction with the separate Vegetation and Wildlife Habitat Mapping Study and the Evaluation of Wildlife Habitat Use Study (see Sections 11.5 and 10.19, respectively), and to estimate the number of migrating and breeding waterbirds that may be affected by the Project.

### **10.15.2. Existing Information and Need for Additional Information**

Existing information on the distribution and abundance of waterbirds in the Project area during the breeding and migration seasons is mostly based on studies conducted in 1980 and 1981 for the Alaska Power Authority (APA) Susitna Hydroelectric Project (Kessel et al. 1982). Data from those studies were used to quantify the level of use of water bodies by migrating and breeding waterbirds. A relative “importance value” was determined for each water body surveyed in each migration season, incorporating the number of species, the number of birds, and the density of birds found on the water body in relation to the overall numbers and densities recorded on the surveys (Kessel et al. 1982). Those study results provide a good knowledge base concerning waterbird use of the Project area three decades ago; however, because the population numbers of numerous species have changed in the past 30 years, new waterbird surveys are needed to elucidate the current distribution and abundance of breeding and migrating waterbirds in the Project area.

More recent survey data on breeding waterbirds in the upper Susitna River basin has been collected annually during U.S. Fish and Wildlife Service (USFWS) waterfowl breeding population surveys (Mallek and Groves 2011a), but only a few transects of the Stratum 2–Nelchina survey area (Mallek and Groves 2011b) are located in the Project area. Those transects occur east of the proposed reservoir near the Oshetna River, an area of low topographic relief where the density of lakes, ponds, and wetlands is relatively high.

The population of Trumpeter Swans is an example of a waterbird species whose population has changed substantially in the last 30 years (Conant et al. 2007). A complete census of Trumpeter Swans on their breeding grounds in Alaska began in 1968 and was repeated at 5-year intervals between 1975 and 2005 (Conant et al. 2007). Together, two survey areas (Unit 3–Gulkana and Unit 5–Cook Inlet) include the entire Susitna River basin (Conant et al. 2007). The population of Trumpeter Swans summering in Alaska has increased since 1975 and breeding has expanded into peripheral habitat. No census was conducted in 2010, so information on the distribution and abundance of Trumpeter Swans in the Project area is out of date and new surveys are needed.

Waterbird productivity was evaluated in 1981 using ground surveys of water bodies within proposed impoundment areas and access routes associated with the APA Susitna Hydroelectric Project. Those surveys provide historical data for the area 30 years ago, but need to be updated. Current surveys addressing waterbird productivity need to be conducted in areas of proposed facility locations, road and transmission corridors, and any areas affected by the Project within and near the inundation zone.

No current information exists on the distribution and abundance of Harlequin Ducks in the middle and upper Susitna River basin. The Harlequin Duck is a species of conservation concern that nests and raises broods almost exclusively in mountain stream drainages. New surveys need to be conducted to assess the distribution and abundance of Harlequin Ducks breeding in the Project area.

### **10.15.3. Study Area**

The study area for waterbirds will encompass lakes, ponds, rivers, streams, and flooded wetlands within a 3-mile buffer area around the Project area (Figure 10.15-1). The 3-mile buffer includes nearly all of the 65 water bodies surveyed for the original APA Susitna Hydroelectric Project in

the 1980s (Kessel et al. 1982), most of which occur in relatively discrete groupings (e.g., see Pre-Application Document [PAD] Figure 4.6-16; AEA 2011). The study area boundary has been extended farther than 3 miles in several places to include water bodies surveyed by Kessel et al. (1982), such as Stephan Lake, Clarence Lake, and other unnamed water bodies south of the Susitna River between Kosina Creek and the Oshetna River, but six large lakes surveyed (Kessel's numbers 131–136) between the mouths of the Tyone and Maclaren rivers will be omitted because they are located well upstream from the area that may be affected by the Project.

Rather than specifying a minimum water body size to be surveyed for the lake-to-lake surveys, the most efficient flight path through each water body group, and linking to other water body groups, will be determined by an experienced waterbird biologist before the surveys begin, to maximize the number of water bodies covered. That same route will be repeated on each migration and breeding-pair survey using Global Positioning System (GPS) navigation; brood surveys will concentrate on the subset of those water bodies located within 1 mile around the locations and alignments of proposed Project infrastructure, including access road and transmission corridors. The survey route will be developed by reviewing U.S. Geological Survey (USGS) 1:63,360-scale topographic maps and high-resolution aerial or satellite imagery, as available, and a GPS route file will be created for navigational use during the survey. It is anticipated that all water bodies 2 hectares (5 acres) or more in size will be surveyed, as well as many smaller ponds located between those larger water bodies. This approach will provide more complete survey coverage than selecting a random sample from all water bodies in the study area. All water bodies sampled will be entered into a Geographic Information System (GIS) database to permit measurement of their area for use in calculating waterbird density.

All rivers and streams flowing through the study area buffer will be surveyed for breeding Harlequin Ducks. These stream surveys will extend outside the 3-mile study-area buffer where necessary to cover suitable habitats farther upstream. Habitat suitability will be evaluated during the first pre-nesting survey for use in planning the three subsequent surveys.

A rectangular area has been delineated east of the upper end of the reservoir inundation zone ("transect block" in Figure 10.5-1) in an area of low topographic relief with a high density of water bodies. The transect block will be sampled during breeding-pair surveys using a transect sampling approach, rather than attempting to cover all of the water bodies completely in a lake-to-lake pattern.

#### **10.15.4. Study Methods**

##### *10.15.4.1. Spring and Fall Migration*

###### *10.15.4.1.1. Aerial Surveys*

Waterbirds use a broad range of lakes, ponds, rivers, and flooded wetlands throughout the Project area during migration. The most effective means of assessing the distribution and abundance of waterbirds over such a large area is aerial survey. Because of the distribution of water bodies in relatively discrete, irregularly spaced groupings in most of the study area, a lake-to-lake survey pattern is the most efficient survey approach, in which each lake is circled to count waterbirds in the water and on the shore. Waterbirds often use rivers and streams for staging during early spring when lakes are covered by ice, so surveys will be flown parallel to river and stream courses.

Aerial surveys of waterbirds in Alaska typically are conducted using either a fixed-wing aircraft or a helicopter, with the choice of platform depending, in part, on the topography of the survey area. Because of the canyons and mountainous terrain in the Project area, a small piston helicopter (Robinson R-44) is the preferred waterbird survey platform to ensure good visibility, survey efficiency, and safety in maneuvering.

To adequately characterize the period of migration and avoid missing migration peaks for various species of waterbirds, surveys will be conducted at 5-day intervals during the spring (late April to early June) and fall (mid-August to mid-October) migration periods, resulting in 10–11 surveys in spring and 13–14 surveys in fall, weather permitting. Each survey is expected to take approximately 2 days to complete. The spring migration surveys will transition directly into the breeding-pair surveys with no break in timing, as is described below (Section 10.15.4.2.1).

A single, experienced observer will record all data on a hand-held digital recorder, which will be transcribed later into a computer database for analysis. Data will be summarized by species, species group, lake group or river segment, date of survey, and survey area. The survey results will be used to evaluate species composition and the timing of migration and to identify water bodies important to migrating waterbirds. Flight lines will be recorded on each survey using a GPS receiver.

#### *10.15.4.1.2. Migration Study*

To acquire current information on the volume and flight directions of birds migrating through the study area, an intensive study of bird migration will be conducted using a combination of visual surveys and radar monitoring. The sampling site for the migration study will be established on the benchland just northeast of the proposed dam site, in the vicinity of the proposed Project camp. Although this study component is described here in the Waterbird Migration, Breeding, and Habitat Use Study plan, it is important to note that the sampling design will also provide data for the Landbird and Shorebird Migration, Breeding, and Habitat Use Study (Section 10.16).

Diurnal visual observations will be conducted during daylight hours (sunrise to sundown) from late April to early June and from mid-August to mid-October. Using binoculars and spotting scopes, observers will record data along four visual transect lines (oriented in the cardinal directions—north, east, south, west) during 25-minute sampling sessions, separated by 5-minute break periods during which weather data will be recorded. Data recorded for each bird observation will include date, time, species (or taxon), flock size, transect crossed, distance crossed (distance from observer), flight direction, flight behavior, minimal flight-altitude category, and an estimate of actual minimal flight altitude.

A portable marine radar that functions in both surveillance and vertical modes will be set up at the sampling site and will be powered by a portable generator. The radar (Furuno Model FR-1510 MKIII; Furuno Electric Company, Nishinomiya, Japan) is a standard X-band marine radar transmitting at 9.410 GHz through a 2-m-long slotted wave guide (antenna) with a peak power output of 12 kW. The antenna has a beam width of 1.23° (horizontal) × 25° (vertical) and a side lobe of ±10–20°. Range accuracy is 1% of the maximal range of the scale in use or 30 m (whichever is greater) and bearing accuracy is ±1°. This radar can be operated at a variety of ranges (0.5–133 km) and pulse lengths (0.07–1.0 µsec). A pulse length of 0.07 µsec will be used while operating at the 1.5-km range to sample the flight activity of small-bodied birds (e.g., songbirds). A longer pulse length (0.3 µsec) will be used while operating at the 6-km range to

sample the flight activity of large-bodied birds (e.g., waterfowl, cranes, raptors). At shorter pulse lengths, echo resolution is improved (giving more accurate information on target identification, location, and distance); whereas, at longer pulse lengths, echo detection is improved (increasing the probability of detecting a target). An echo is a picture of a target on the radar monitor; a target is one or more birds (or bats) that are flying so closely together that the radar displays them as one echo on the display monitor. The radar has a digital color display with several useful features, including true north correction for the display screen (to determine flight directions), color-coded echoes (to differentiate the strength of return signals), and on-screen plotting of a sequence of echoes (to depict flight paths). Because targets are plotted with every sweep of the antenna (i.e., every 2.5 sec) and because ground speed is directly proportional to the distance between consecutive echoes, ground speeds of plotted targets can be estimated to the nearest 5 km/h with a hand-held scale.

Radar data will be collected in several 1-hour sampling sessions throughout the night (from shortly after sunset to just before sunrise) and diurnal radar sampling sessions will be conducted during the day (shifting 3-hour blocks from morning to evening). Each 1-hour radar sampling session will consist of (1) one 10-min period to collect weather data and adjust the radar to surveillance mode; (2) one 10-min period with the radar in surveillance mode (1.5-km range) for collection of information on migration passage rates or small-bodied birds; (3) one 10-min period with the radar in surveillance mode (1.5-km range) for collection of information on ground speed, flight direction, tangential range (minimal perpendicular distance to the radar laboratory), transect crossed (north, south, east, and west), and the number of individuals (if known) of small-bodied birds; (4) one 10-min period with the radar in surveillance mode (6-km range) for collection of information on both passage rates of large-bodied birds and information on their groundspeed, flight direction, tangential range (minimal perpendicular distance to the radar laboratory), transect crossed (north, south, east, and west), and the number of individuals (if known); (5) one 5-min period to adjust the radar to vertical mode; and (5) one 15-min period with the radar in vertical mode (1.5-km range) to collect information on flight altitudes and flight behavior. All hours of radar data will be recorded using an automated image frame-grabber device (Model VGA2USB, Epiphan Systems Inc., Ottawa, Ontario, Canada) that will enable continuous collection of a record of high-quality lossless radar images, with a resolution identical to that of the radar monitor.

Nocturnal audiovisual surveys will be conducted during the first 2 hours of nocturnal radar sampling in both spring and fall, and will include two sessions of visual sampling separated by short periods during which weather data will be recorded. The sampling period will be adjusted as daylength changes during the migration periods. The observer will use binoculars during crepuscular periods and night-vision goggles during dark hours, aided by spotlights outfitted with infrared filters to illuminate targets flying overhead. For each bird or flock of birds detected visually, the following data will be collected: species or taxon, flight direction, flight altitude, and flight behavior. Weather data recorded before each radar or visual sampling session will include wind direction, average wind speed, cloud cover, ceiling height, light conditions, precipitation, air temperature, and barometric pressure.

Data collected in this study on flight volume, altitudes, and directions among all species and taxa will be compared with data collected in similar studies at Tok in the upper Tanana River valley and Gakona in the Copper River valley (Cooper et al. 1991a, 1991b; Cooper and Ritchie 1995),

the Tanana Flats and Alaska Range foothills near Healy (Day et al. 2007; Shook et al. 2006, 2011), and Fire Island (Day et al. 2005), at minimum.

#### *10.15.4.2. Breeding Season*

##### *10.15.4.2.1. Breeding-pair Surveys*

Two different survey approaches will be used for breeding-pair surveys, depending on the location of the water bodies being surveyed. In most of the study area, the same lake-to-lake survey approach will be used as during the migration surveys, with no break in timing between the spring migration and breeding survey periods. In the designated transect survey block in the easternmost portion of the study area, however, a sampling approach will be used to survey 400-meter-wide strips along transects spaced at 1-mile intervals, providing sample coverage of approximately 25 percent of the survey block.

Surveys for breeding waterbirds, primarily waterfowl, will generally follow standard USFWS protocols (USFWS 1987; USFWS and CWS 1987). The survey lines in the transect block will be aligned to cover the largest possible number of water bodies and wetlands. The placement of the transect lines, which will be oriented systematically along the long axis of the survey block, will be determined before the survey using aerial imagery or topographic maps and GIS.

Breeding-pair surveys are typically conducted in a fixed-wing aircraft; however, it is likely that a small piston helicopter will be used for all aerial surveys of waterbirds in this study. Flight altitude will be low (125–200 feet above ground level, with the lower altitude being used for the transect surveys) to permit observation of birds without having to rely on binoculars, although binoculars will be used where necessary to confirm species identity. In the lake-to-lake surveys, a single observer will record data over the entire area of the water bodies surveyed. In the transect surveys, one observer will search for waterbirds in a 200-meter (656-foot) swath on each side of the aircraft (total of two observers searching a total strip width of 400 meters, or 1,312 feet) while the pilot navigates the transect lines using a GPS receiver. Observations will be recorded on hand-held digital voice recorders for later transcription and transfer to a computer database for analysis. Survey data will be used to calculate the estimated densities of each species of waterfowl and identify areas important to breeding waterfowl.

The timing of the breeding-pair surveys will be determined by evaluating the chronology of break-up and spring melt conditions each year, which will be monitored throughout the spring migration surveys. Breeding-pair surveys typically are flown in late May or early June, depending on location and elevation, when pairs are present on territories but females are not yet spending time on nests. Survey timing can affect results because the nesting phenology of dabbling ducks is earlier than that of diving ducks, and some dabbling duck species can be missed if the survey occurs too late, after the cryptically colored females are on nests and the more brightly colored males have left the area. Two surveys, spaced about 10 days apart, will be conducted to target the expected peaks of breeding for dabbling and diving ducks. The two breeding-pair surveys will be timed to coincide with the peak presence of pairs and males of dabbling ducks and diving ducks, respectively, to account for the differences in migration timing and turnover of those two general categories of ducks. Each survey is expected to take approximately two days to complete. Weather and visibility conditions will be recorded during surveys to assess the quality of the information recorded, and surveys will not proceed unless conditions are suitable. Standard protocols (USFWS and CWS 1987, Smith 1995) will be

followed to convert raw survey counts to indicated total population indices and species-specific correction factors will be applied to the indices to derive population estimates of each species detected in the transect strips for which correction factors are available.

#### *10.15.4.2.2. Harlequin Duck Surveys*

In inland areas of Alaska, Harlequin Ducks predominantly use mountain streams for foraging and nest in adjacent shoreline habitats. Male Harlequin Ducks are only present on breeding streams during a short period in spring while courting females. Accordingly, pre-nesting surveys to quantify the number of nesting pairs occupying a stream must be conducted in that short timing window. After hatching, successful females are visible on streams with their broods, and failed breeders often group together.

Surveys for pre-nesting and brood-rearing Harlequin Ducks will be flown in a helicopter with two observers seated on the same side. Surveys generally will be flown in an upriver direction with the helicopter positioned over the bank to provide an unobstructed view of the entire width of the watercourse. Each survey is expected to take approximately two days to complete. Surveys will follow the entire length of tributary streams where suitable nesting habitat is present, even if it means flying outside of the 3-mile study area buffer. The extent of suitable habitat will be defined during the first pre-nesting survey. Observations will be recorded on hand-held digital recorders and with GPS waypoints for later transcription and transfer to a computer database for analysis. Survey data will be used to calculate linear densities (ducks per kilometer) and to identify streams used by breeding Harlequin Ducks.

To account for annual variability in the occurrence of peak numbers of breeding pairs and brood-rearing females on a stream, two years of pre-nesting and brood-rearing surveys will be conducted. Two pre-nesting surveys, spaced 7–10 days apart, will be flown in late May–early June each year and two brood-rearing surveys, spaced 7–10 days apart, will be conducted in late July–early August each year, with the exact timing to be determined using information on environmental conditions and breeding phenology each year.

#### *10.15.4.2.3. Brood Surveys*

Information on waterbirds breeding in specific areas that may be affected by Project infrastructure or activities will be collected by biologists conducting helicopter surveys of suitable lakes, ponds, streams, and flooded wetland complexes. As with the other waterbird surveys, the platform of choice will be a small piston helicopter. These surveys will be conducted at least twice during the brood-rearing period, with the first survey occurring in mid-July and the second approximately a week later, to record the presence of adults accompanied by broods of juveniles. A third survey may be flown, depending on the developmental stages of juveniles observed on the second brood survey. The brood surveys will focus on the water body groups within 1 mile around the locations and alignments of proposed Project infrastructure, excluding portions of the study area located farther than 1 mile away.

Two observers will circumnavigate water bodies in a small helicopter to search for waterbirds with broods. All waterbirds seen will be recorded on field data sheets and brood ages for waterfowl (primarily ducks) will be estimated by classifying them into one of seven age classes based on chick plumage patterns. Survey data will be used to calculate densities of broods and to estimate nest-initiation dates by back-dating (subtracting the estimated age of young and the

average incubation period). Any nest locations of piscivorous waterbirds will be recorded for collection of feathers for analysis of baseline mercury levels for the Mercury Assessment and Potential for Bioaccumulation Study (Section 5.7).

#### *10.15.4.3. Information for Mercury Assessment*

To provide tissue samples for laboratory analysis of mercury levels, feathers of piscivorous waterbirds (e.g., loons, grebes, mergansers, terns) will be collected, after the nesting season, from any nests of those species that are located during aerial surveys for this study or surveys for other Project studies. It is anticipated that some nests of loons, grebes, and Arctic Terns will be found because of their selection of nest sites on islands or shorelines, but merganser nests are unlikely to be found because they nest in tree cavities that would be difficult to locate without radio telemetry. Therefore, to supplement the collection of feathers from nests, prey remains will be collected from nest sites of Peregrine Falcons located in or near the study area. Peregrine Falcons are predators of a variety of birds, including waterbirds, and examination of prey remains is a commonly used technique to investigate their food habits. A permit will be required from the USFWS to collect any migratory bird parts.

In addition to collection of feather samples for laboratory analysis, the scientific literature will be reviewed to compile and synthesize information on the food habits and diets of piscivorous waterbirds in freshwater aquatic systems to support the risk assessment being conducted as part of the Mercury Assessment and Potential for Bioaccumulation Study (Section 5.7). This information was recommended by USFWS in comments on the Pre-Application Document for the Project (letter from USFWS to AEA dated May 31, 2012).

#### *10.15.4.4. Reporting and Deliverables*

Study products will include the following:

- **Electronic copies of field data.** A geospatially referenced relational database will be developed that incorporates all historic and current data, including nesting and brood-rearing locations for each species. Naming conventions of files and data fields, spatial resolution, map projections, and metadata descriptions will meet the data standards to be established for the Project.
- **Study Reports.** The Initial Study Report will be completed in February 2014 and the Updated Study Report will be completed in February 2015. The Updated Study Report will summarize the results for both years.

#### **10.15.5. Consistency with Generally Accepted Scientific Practice**

The Waterbird Migration, Breeding, and Habitat Use Study will be conducted using standard waterfowl aerial survey techniques, including those described in the current USFWS *Standard Operating Procedures for Aerial Waterfowl Breeding Ground Population and Habitat Surveys in North America* (USFWS and CWS 1987). These same techniques have been used successfully to survey for migrant and breeding waterbirds on other large-scale projects in Alaska (e.g., PLP 2011). Stream surveys of Harlequin Ducks by helicopter have been used effectively in numerous studies in Alaska and Canada (McCaffery and Harwood 1994; Morgart 1998; Kneteman and Hubbs 2000; Paton 2000; Savage 2000; MacDonald 2003; PLP 2011). The diurnal visual and nocturnal radar techniques proposed in this plan have become standard methods for studying bird



migration (e.g., Gauthreaux 1980; Cooper et al. 1991a, 1991b; Gauthreaux and Belser 2003), including in Alaska (Cooper et al. 1991a, 1991b; Cooper and Ritchie 1995; Day et al. 2005, 2007; Shook et al. 2006, 2011).

#### **10.15.6. Schedule**

The migration study using coordinated radar and visual sampling is proposed for the spring (late April–early June) and fall (mid-August–mid-October) migration periods in 2013, with the decision to conduct a second year of sampling being dependent on the results obtained in 2013.

For the other study components described above, the same seasonal schedule will be followed in both 2013 and 2014 (Table 10.15-1). The timing of some surveys, particularly in spring and summer, will depend on ice break-up and the nesting phenology for the year. Approximately 10–11 spring migration/breeding-pair surveys will be flown between late April (start date determined by the timing of river break-up and lake moat formation each year) and mid-June at intervals of 5 days. Two breeding-pair transect surveys will be flown in the first half of June. At least two brood surveys (possibly three, depending on the developmental stages of juveniles on the second survey) will be flown beginning in mid-July each year. Two pre-nesting surveys for Harlequin Ducks will be flown in late May/early June and two brood-rearing surveys will be flown in late July/early August. Fall migration surveys will begin in mid-August and will continue until mid-October at intervals of 5 days. After each aerial survey, data will be transcribed, reviewed, and entered into a database for final quality assurance/quality control (QA/QC) review. Data analysis will be conducted progressively throughout summer and fall to allow for rapid completion in November and December. The Initial Study Report will be completed by February 2014 and the Updated Study Report will be completed by February 2015. Study updates will be provided during Technical Workgroup meetings, which will be held quarterly during 2013 and 2014.

#### **10.15.7. Relationship with Other Studies**

Except for current information each spring about river break-up conditions from the study of Ice Processes in the Susitna River (Section 7.6), which will be used to help determine the start date of spring migration surveys, the Waterbird Migration, Breeding, and Habitat Use Study will not require specific information inputs from other studies, but will provide information to several other studies (Figure 10.15-2). The various types of surveys conducted for this study will provide information that will be used in the assessment of Project impacts and development of appropriate protection, mitigation, and enhancement (PM&E) measures, which will be conducted in 2015 for the FERC License Application. The ground-based visual and radar surveys during spring and fall migrations in 2013 will be conducted concurrently with the Landbird and Shorebird Migration, Breeding, and Habitat Use Study (Section 10.16) to document the nature of migratory movements of a broad variety of birds near the proposed dam and associated facilities. Information regarding habitat use and abundance among different habitats will be incorporated into the habitat ranking matrix constructed for the Evaluation of Wildlife Habitat Use (Section 10.19), which will be used to assess the potential impacts and to develop PM&E measures, as appropriate, for the FERC License Application. Information on the food habits and diets of piscivorous waterbirds will be synthesized for use in the Mercury Assessment and Potential for Bioaccumulation Study (Section 5.7), and nest locations of waterbird species found during breeding-season surveys will be recorded to enable collection of feather samples (after the nests

are vacated) for baseline characterization of mercury levels by the Mercury Assessment and Potential for Bioaccumulation Study (Section 5.7).

The primary impact mechanisms of the Project on waterbirds may include the following:

- Permanent direct and indirect habitat loss and alteration.
- Temporary direct and indirect habitat loss and alteration.
- Direct behavioral impacts, such as attraction or avoidance, resulting from vehicular use, noise, and increased human presence associated with Project construction or operation.
- Indirect behavioral impacts to wildlife, such as attraction or avoidance, resulting from changes in vehicular use, noise, and increased human presence associated with increased subsistence or recreational access that may be facilitated by Project development.
- Mortality due to increased subsistence and recreational harvest that may be facilitated by improvements in human access that result from Project development.
- Changes in mortality due to predation that may result from changes in the abundance and distribution of waterfowl predators, including both mammalian and avian carnivores.
- Direct mortality due to strikes with vehicles, power lines, towers, or other Project facilities; exposure to contaminants; and attraction to garbage and human activity.

Impacts associated with habitat loss and alteration, attraction and avoidance, and direct mortality will occur primarily in the Project area, including the impoundment area, access and transmission corridors, and other facility footprints. Impacts associated with increased harvest and changes in predator abundance may occur over a larger area in which changes in both competing mammalian predators and prey species abundance may occur.

During the impact assessment that will be conducted in 2015 for the FERC License Application, data on the distribution, abundance, and productivity of and habitat use by waterbirds in the study area will be used to assess Project impacts on these populations. Impacts of direct and indirect habitat loss and alteration can be assessed through geospatial analysis. When plotted on the wildlife habitat map developed by the Vegetation and Wildlife Habitat Mapping Study in the Upper and Middle Susitna Basin (Section 11.5), the locations of breeding pairs, brood-rearing groups, and staging birds will allow identification of high-value seasonal habitats for each waterbird species. Using GIS software, the direct and indirect impacts of the Project can be evaluated for each waterbird species by overlaying the reservoir impoundment, related infrastructure areas, and access road and power transmission corridors onto the habitat map to calculate loss of preferred or critical habitats.

Additional indirect impacts of habitat loss and alteration and behavioral reactions (such as avoidance) can be estimated by applying various buffer distances, as determined from the literature on the effects of similar projects. In this way, the GIS analysis will be combined with information from the literature to estimate the geographic extent, frequency, duration, and magnitude of Project effects on waterbird populations. Density estimates for breeding pairs and brood-rearing waterbirds in each habitat and linear densities of Harlequin Ducks can be used to estimate the number of birds potentially affected by habitat loss and alteration and by behavioral reactions that may result in avoidance. Location data for each species can be used to assess risks from power lines and other bird strikes for various Project alternatives. Industry standards and best practices (e.g., APLIC and USFWS 2005; APLIC 2006) and other literature reviews (e.g., Evans et al. 2007; Longcore et al. 2008; Gehring et al. 2009) will be consulted when designing and siting transmission lines and lighting Project facilities, to reduce the risk of attraction and

collision of birds with Project infrastructure, while still complying with other federal and state requirements for facility lighting for aviation safety. Any necessary PM&E measures will be developed, as appropriate, by examining the distribution and abundance of species among habitats in relation to the geographical extent and seasonal timing of various Project activities.

#### **10.15.8. Level of Effort and Cost**

The bulk of the costs associated with this study are for the field sampling, data analysis, and reporting; helicopter support costs have not been included in this estimate. The aerial survey portion of the study will require an estimated minimum of 86 person-days, not including weather delays or changes in study design, as indicated below:

- Migration surveys = 20 person-days in spring; 26 person-days in fall
- Breeding-pair surveys = 12 person-days (assuming two surveys per year)
- Harlequin Duck pre-nesting surveys = 8 person-days
- Harlequin Duck brood-rearing surveys = 8 person-days
- Waterbird brood-rearing surveys = 12 person-days

The ground-based migration study employing diurnal visual and nocturnal radar and audiovisual sampling (which also will provide data for the Landbird and Shorebird Migration, Breeding, and Habitat Use Study, Section 10.16) will require a crew of four biologists working day and night shifts over a period of up to 120 days (total for spring and fall fieldwork) and is estimated to cost approximately \$600,000 in 2013. The projected cost of the aerial-survey portion of the waterbird study is approximately \$300,000 annually. Hence, the total cost is estimated at \$900,000 in 2013 and \$300,000 in 2014, for a 2-year total of \$1.2 million. The decision to continue the ground-based migration monitoring task in 2014 will be based on the results obtained in 2013.

#### **10.15.9. Literature Cited**

- AEA. 2011. Pre-Application Document: Susitna–Watana Hydroelectric Project FERC Project No. 14241. December 2011. Prepared for the Federal Energy Regulatory Commission by the Alaska Energy Authority, Anchorage, Alaska.
- APLIC (Avian Power Line Interaction Committee) and USFWS (U.S. Fish and Wildlife Service). 2005. Avian Protection Plan (APP) guidelines. Prepared by the Edison Electric Institute’s Avian Power Line Interaction Committee and U.S. Fish and Wildlife Service.
- APLIC. 2006. Suggested practices for avian protection on power lines: the state of the art in 2006. Edison Electric Institute, APLIC, and the California Energy Commission, Washington, DC, and Sacramento, CA.
- Conant, B., J. I. Hodges, D. J. Groves, and J. G. King. 2007. Alaska Trumpeter Swan status report, 2005. U. S. Fish and Wildlife Service, Juneau. 49 pp.
- Cooper, B. A., and R. J. Ritchie. 1995. The altitude of bird migration in east-central Alaska: a radar and visual study. *Journal of Field Ornithology* 66: 590–608.
- Cooper, B. A., R. H. Day, R. J. Ritchie, and C. L. Cranor. 1991a. An improved marine radar system for studies of bird migration. *Journal of Field Ornithology* 62: 367–377.

- Cooper, B. A., R. J. Ritchie, B. A. Anderson, and L. C. Byrne. 1991b. Alaska Over-the-Horizon Backscatter Radar System: a synthesis of the Avian Research Program, 1987–1990. Report to the Arctic Environmental Information and Data Center, Anchorage, and the U.S. Department of the Air Force, Hanscom Air Force Base, Massachusetts, by Alaska Biological Research, Inc., Fairbanks. 309 pp.
- Day, R. H., R. J. Ritchie, J. R. Rose, and A. M. Wildman. 2007. Avian studies along the proposed Northern Intertie powerline project, 1999–2000 (2007 compilation). Report prepared for Golden Valley Electric Association, Inc., Fairbanks, by ABR, Inc.—Environmental Research & Services, Fairbanks. 172 pp.
- Day, R. H., R. J. Ritchie, J. R. Rose, and G. V. Frost. 2005. Bird migration near Fire Island, Cook Inlet, Alaska, spring and fall 2004. Report prepared for Chugach Electric Association, Inc., Anchorage, by ABR, Inc.—Environmental Research & Services, Fairbanks. 128 pp.
- Evans, W. R., Y. Akashi, N. S. Altman, and A. M. Manville, Jr. 2007. Response of night-migrating birds in cloud to colored and flashing light. Report to the Communications Tower Working Group. 25 pp.
- Gauthreaux, S. A. 1980. *Direct visual and radar methods for the detection, quantification, and prediction of bird migration*. Clemson University, Clemson, SC. 67 pp.
- Gauthreaux, S. A., and C. G. Belser. 2003. Radar ornithology and biological conservation. *Auk* 120: 266–277.
- Gehring, J., P. Kerlinger, and A. M. Manville, Jr. 2009. Communication towers, lights, and birds: successful methods of reducing the frequency of avian collisions. *Ecological Applications* 19: 505–514.
- Kessel, B., S. O. MacDonald, D. D. Gibson, B. A. Cooper, and B. A. Anderson. 1982. Susitna Hydroelectric Project environmental studies, Phase I final report—Subtask 7.11: Birds and non-game mammals. Report by University of Alaska Museum, Fairbanks, and Terrestrial Environmental Specialists, Inc., Phoenix, NY for Alaska Power Authority, Anchorage. 149 pp.
- Kneteman, J., and A. Hubbs. 2000. Harlequin Duck monitoring in the northern east slopes of Alberta: 1998–2000 preliminary results. Alberta Sustainable Resource Development. Fisheries and Wildlife Management Division, Alberta Species at Risk Report No. 11, Edmonton, AB. 18 pp.
- Longcore, T., C. Rich, and S. A. Gauthreaux, Jr.. 2008. Height, guy wires, and steady-burning lights increase hazards of communication towers to nocturnal migrants: a review and meta-analysis. *The Auk* 125: 485–492.
- MacDonald, R. 2003. Harlequin Duck breeding pair and brood surveys, Togiak National Wildlife Refuge, Alaska, 1998–2002. U.S. Fish and Wildlife Service, Togiak National Wildlife Refuge, Dillingham, AK. 40 pp.
- Mallek, E. J., and D. J. Groves. 2011a. Alaska–Yukon waterfowl breeding population survey. U.S. Fish and Wildlife Service, Fairbanks and Juneau, AK. 30 pp.

- Mallek, E. J., and D. J. Groves. 2011b. Map of transects within Stratum 2 (Nelchina) of the Alaska–Yukon Waterfowl Breeding Population Survey [map of Interior Alaska portion of the survey]. 1 p.
- McCaffery, B. J., and C. M. Harwood. 1994. Distribution and relative abundance of Harlequin Ducks in the southwest Kuskokwim Mountains. U.S. Fish and Wildlife Service, Yukon Delta National Wildlife Refuge, Bethel, AK. 13 pp.
- Morgart, J. R. 1998. Kilbuck Mountains Harlequin Duck breeding-pair survey. U.S. Fish and Wildlife Service, Yukon Delta National Wildlife Refuge, Bethel, AK. 18 pp.
- Paton, D. 2000. Harlequin Duck surveys of Oldman River Basin, 2000. Alberta Sustainable Resource Development. Fisheries and Wildlife Management Division, Alberta Species at Risk Report No. 20, Edmonton, AB. 37 pp.
- PLP (Pebble Limited Partnership). 2011. Pebble Project Environmental Baseline Document, 2004 through 2008. Pebble Limited Partnership, Anchorage, AK. Available online: <http://www.pebbleresearch.com/> (accessed June 16, 2012).
- Savage, S. 2000. Harlequin Duck stream survey, Alaska Peninsula/Becharof National Wildlife Refuge, Alaska, July 2000. U.S. Fish and Wildlife Service, Alaska Peninsula/Becharof National Wildlife Refuge Complex, King Salmon, AK. 36 pp.
- Shook, J. E., R. H. Day, and R. J. Ritchie. 2006. Monitoring interactions of birds with the Northern Intertie powerline, Interior Alaska, spring and fall 2004–2005. Report prepared for Golden Valley Electric Association, Inc., Fairbanks, by ABR, Inc.—Environmental Research & Services, Fairbanks. 46 pp.
- Shook, J. E., J. H. Plissner, L. B. Attanas, R. H. Day, and R. J. Ritchie. 2011. Avian migration studies at the proposed Eva Creek windfarm: spring and fall migration 2010. Report prepared for Golden Valley Electric Association, Inc., Fairbanks, by ABR, Inc.—Environmental Research & Services, Fairbanks. 79 pp.
- Smith, G. W. 1995. A critical review of the aerial and ground surveys of the breeding waterfowl surveys in North America. U.S. Fish and Wildlife Service, Biological Science Report No. 5.
- USFWS (U.S. Fish and Wildlife Service). 1987. Trumpeter and Tundra Swan survey protocol update. U. S. Fish and Wildlife Service Memorandum, Juneau, AK.
- USFWS and CWS (Environment Canada, Canadian Wildlife Service). 1987. *Standard operating procedures for aerial breeding-ground population and habitat surveys in North America*. Migratory Bird and Habitat Research Laboratory, Patuxent Wildlife Research Center, Laurel, MD.

### 10.15.10. Tables

**Table 10.15-1. Schedule for implementation of the Waterbird Migration, Breeding, and Habitat Use Study.**

Activity	2013				2014				2015
	1 Q	2 Q	3 Q	4 Q	1 Q	2 Q	3 Q	4 Q	1 Q
Spring migration surveys (ground-based visual & radar monitoring)		—				—			
Spring migration aerial surveys, transitioning to breeding-pair surveys; Breeding-pair transect surveys; Pre-nesting surveys for Harlequin Ducks		—				—			
Brood surveys in water bodies; Harlequin Duck brood surveys in streams			—				—		
Fall migration surveys (ground-based visual & radar monitoring)			—	—			—	—	
Fall migration aerial surveys			—	—			—	—	
Data QA/QC and analyses			—	—	—		—	—	—
Initial Study Report				—	△				
Updated Study Report								—	▲

Legend:

- Planned Activity
- △ Initial Study Report
- ▲ Updated Study Report

10.15.11. Figures




Figure 10.15-1. Waterbird Migration, Breeding, and Habitat Use Study area.

**STUDY INTERDEPENDENCIES FOR WATERBIRD STUDY**




Figure 10.15-2. Study interdependencies for the waterbird study.



## **10.16. Landbird and Shorebird Migration, Breeding, and Habitat Use Study**

### **10.16.1. General Description of the Proposed Study**

The Landbird and Shorebird Migration, Breeding, and Habitat Use Study is planned as a two-year study (2013–2014). Results from the first year of work in 2013 will be used to update existing information and to fine-tune the field survey methods and survey areas in 2014, if necessary. The landbird and shorebird study will employ three basic survey methods: ground-based point-count surveys for breeding birds, a boat-based survey for colonially nesting swallows, and ground-based monitoring of migration using a combination of daytime visual observations and nocturnal radar sampling. The point-count surveys are intended to record all birds seen or heard and to estimate the distance to each bird detected. Point-count surveys, which were designed to count singing male passerines, are now the preferred method for inventory and monitoring efforts for landbirds in remote, roadless terrain in Alaska (Handel and Cady 2004; ALMS 2010). These methods also have been adopted for shorebirds (ASG 2008) and are especially appropriate in forested landscapes, where shorebirds typically occur in low densities and where plot-based methods would yield few observations, even with a relatively large survey effort. The survey of colonially nesting swallows will focus on suitable habitats within the proposed reservoir inundation zone. The boat-based survey will identify swallow nesting colonies and potential nesting habitat.

#### **Study Goal and Objectives**

The goal of this study is to collect baseline data on the occurrence and habitat use of breeding landbirds and shorebirds in the Project area to enable assessments of the direct, indirect, and cumulative impacts on these birds from construction and operation of the proposed Project. This study will include species of conservation concern, both landbirds and shorebirds, that are known or expected to occur in the Project area (see AEA 2011), as well as numerous other species that are protected under the federal Migratory Bird Treaty Act (see Section 10.3).

The study has four specific objectives:

- Collect data on the distribution and abundance of landbirds and shorebirds during the summer breeding season.
- Identify habitat associations for landbirds and shorebirds.
- Evaluate changes in distribution, abundance, and habitat use of landbirds and shorebirds through comparison with historical data.
- Characterize the timing, volume, direction, and altitude of landbirds and shorebirds migrating through the dam and camp facilities area.

To achieve these objectives, the following surveys and analyses will be required:

- Conduct ground-based point-count surveys to collect field data on the distribution and abundance of landbirds and shorebirds in the study area during the summer breeding season.
- Conduct focused point-count and linear walking surveys in riparian and lacustrine habitats, targeting piscivorous species and other species typical of fluvial, riparian, and lacustrine habitats, which often are under-represented in standard point-count surveys.

- Conduct boat-based surveys of colonially nesting swallows in riparian habitats within the reservoir inundation zone.
- Collect habitat-use data for landbirds and shorebirds during the point-count surveys to inform the Evaluation of Wildlife Habitat Use Study (Section 10.19), which will be the first step in quantifying habitat change (i.e., gain/loss and alteration) for landbirds and shorebirds from the proposed Project.
- Review the literature on the foraging habits and diets of piscivorous and partly piscivorous landbird and shorebird species (e.g., Belted Kingfisher, American Dipper, Spotted Sandpiper), which will be used to inform the Mercury Assessment and Potential for Bioaccumulation Study (see Section 5.7).
- Conduct visual migration-watch surveys and radar sampling in the immediate vicinity of the dam, powerhouse, and camp facilities.
- Review historical (Alaska Power Authority [APA] Susitna Hydroelectric Project) data on landbirds and shorebirds for comparison with the current data from this study, to evaluate any changes in distribution, abundance, and habitat use over the intervening 30 years. Many species of migratory birds have suffered population declines in recent decades, so these comparisons may also provide information on the population status of those species in the Project area.

#### **10.16.2. Existing Information and Need for Additional Information**

In 1980 and 1981, breeding landbirds and some shorebirds were surveyed for the APA Susitna Hydroelectric Project using modified territory-mapping methods, which involved repeated visits between May 20 and July 3 to 12 study plots, each 10 hectares (24.7 acres) in size (Kessel et al. 1982; AEA 2011). Each plot was placed in an area of homogeneous habitat, as defined using Kessel's avian habitat classification (Kessel 1979). At that time, territory mapping was the standard method for surveying landbirds. Because each plot was surveyed repeatedly, substantial information on bird occurrence and habitat use was obtained for the limited area encompassed by those 12 plots. However, because only 12 plots were sampled in homogeneous habitats, the data did not adequately address spatial variability in bird occurrence and habitat use across the broader study area. Some additional information on shorebird occurrence was obtained during ground-based surveys of lakes, ponds, and wetlands for waterbirds (Kessel et al. 1982), but focused surveys for breeding shorebirds were not conducted. No studies of landbirds or shorebirds have been conducted more recently in the Project area (AEA 2011).

During the surveys by Kessel et al. (1982), four species of swallows were observed in the Project area: Tree Swallow, Bank Swallow, Cliff Swallow, and Violet-green Swallow. Violet-green and Tree swallows were considered fairly common, whereas Bank and Cliff swallows were considered uncommon, and all four species were either confirmed or suspected to nest in the study area. A nesting colony of 25 pairs of Bank Swallows was recorded along upper Watana Creek and three colonies of Cliff Swallows were observed at Watana and Clarence lakes. The distribution of avian survey plots and incomplete representation of habitat types suitable for swallows likely led to an underestimate of swallow abundance and distribution in the Project area by Kessel et al. (1982). No studies of swallows have been conducted more recently in the study area.

Because of the limitations in extrapolating results from intensive surveys of territory-mapping plots to the larger Project area, it will be necessary to study these species groups again using

currently accepted protocols (point-count surveys), which allow large landscapes to be sampled adequately and which provide more data on variability in habitat use. Because the most recent surveys for landbirds and shorebirds were conducted over 30 years ago, and because populations of these birds and their habitats have likely changed during that period, new studies are recommended. Current data on the distribution, abundance, and habitat use of landbirds and shorebirds is necessary to be able to adequately assess the impacts from the proposed Project on these species.

Point-count surveys are not always effective in riparian habitats, where the auditory capacity of observers can be reduced in some locations because of rapidly flowing water, and are sub-optimal for species such as swallows, that have highly clumped distributions (Swanson and Nigro 2003). Because much of the swallow nesting habitat in the study area is difficult to access or observe on foot, a separate survey effort has been designed to assess the distribution and abundance of colonially nesting swallows in riparian habitats.

No migration surveys for landbirds and shorebirds were conducted specifically for the original APA Susitna Hydroelectric Project studies, although information was compiled opportunistically (Kessel et al. 1982). The U.S. Fish and Wildlife Service (USFWS) is concerned about the potential for attraction of migrating landbirds and shorebirds to artificially lighted facilities constructed for the Project, potentially resulting in mortality from collisions, as expressed in comments in various meetings (see Appendices 3 and 4).

### **10.16.3. Study Area**

The study area for the breeding landbird and shorebird point-count survey includes a 2-mile buffer zone around the proposed reservoir impoundment zone, dam and camp facilities area, access road and transmission corridor alternatives, and material sites (Figure 10.16-1). All direct and indirect effects of the proposed Project on landbirds and shorebirds and their habitats in the upper Susitna basin are expected to be encompassed by this 2-mile buffer.

The point-count and linear walking surveys in riparian and lacustrine habitats will be conducted in the primary riparian and lacustrine habitats in the inundation zone, the infrastructure area surrounding the site of the proposed dam, and in riparian habitats along the Susitna River immediately below the location of the proposed dam.

The survey area for colonially nesting swallows includes riparian habitats along the Susitna River and its tributaries within the reservoir inundation zone. The Susitna River in the inundation zone is an alluvial river located in a shallow canyon and is moderately braided with a low gradient. Tributaries to the Susitna River (e.g., Deadman, Tsusena, Watana, Jay, and Kosina creeks) are lower volume, steeper gradient, clearwater streams with deeply incised canyons along their lower reaches. Habitat features to be examined during the survey will include all riverbanks and cliffs adjacent to these rivers and streams.

The sampling site for the migration study component will be located on the adjacent benchland just above the dam site on the north side of the river (Figure 10.16-1).

## **10.16.4. Study Methods**

### *10.16.4.1. Point-count Surveys*

#### *10.16.4.1.1 Study Design*

The proposed methods for the breeding landbird and shorebird study are ground-based point-count surveys, in which all birds seen or heard are recorded, along with an estimate of the horizontal distance to each bird observed. Point-count surveys, which were designed to count singing male passerine birds, are now the preferred method for inventory and monitoring efforts for landbirds in remote, roadless landscapes in Alaska (Handel and Cady 2004; ALMS 2010). These methods have been adopted for shorebirds (ASG 2008) and are especially appropriate in forested landscapes, where shorebirds typically occur in low densities and where plot-based methods would yield few observations, even with a relatively large survey effort.

Point-count surveys are appropriate for large development projects that affect a large area and can include many different types of habitats. The sample points can be distributed across the landscape and allocated among habitat types to ensure that all prominent habitat types are sampled. Because management agencies in Alaska are increasingly concerned with landbird and shorebird species of conservation concern (which are generally uncommon), and because it is important to sample many different occurrences of each habitat type to detect uncommon species (which are patchy in occurrence across the landscape), this study has been designed so that point-counts are allocated in as many different occurrences of each of the prominent habitat types in the study area. In 2013, point-count sampling locations will be distributed using a pseudo-stratified random plot allocation procedure based on aerial photosignatures as the sampling strata (because a current, complete habitat map will not be available by spring 2013). The plot allocation methods may change in 2014 after a current, complete habitat map is available. This procedure will result in adequate sampling of habitats, over two years of surveys, so that habitat-use evaluations for landbirds and shorebirds will be supported by Project area-specific data. These habitat-use evaluations (see Section 10.19) are a critical link in conducting quantitative assessments of habitat loss and alteration for breeding landbirds and shorebirds.

Several species of landbirds and shorebirds are not commonly recorded in standard point-count surveys allocated randomly across available habitats, but are known to be closely associated with riparian and lacustrine habitats (e.g., Belted Kingfisher, American Dipper, Semipalmated Plover, Solitary Sandpiper, Spotted Sandpiper, Wandering Tattler). Therefore, an additional set of point-count surveys will be conducted specifically in riparian and lacustrine habitats that are expected to be affected by Project development (see Section 10.16.4.2 below). These additional surveys were requested by USFWS (see Appendices 3 and 4).

The landbird and shorebird study will be coordinated with the other wildlife studies being conducted for the Project, especially the raptor and waterbird studies, so that sightings of bird species that apply to other studies can inform the qualitative results and reporting efforts among studies.

#### *10.16.4.1.2 Field Surveys*

Point-count field surveys will follow standard protocols for point-counts in Alaska (Handel and Cady 2004; ALMS 2010). These protocols are based on the variable circular-plot point-count

methods and temporally stratified observation periods (Ralph et al. 1995; Buckland et al. 2001; Farnsworth et al. 2002; Rosenstock et al. 2002). Surveys will be conducted during early morning hours to maximize the detection of breeding species, especially singing male passerines. Standard 10-minute observation periods will be used and, to facilitate the collection of habitat-use data, the specific habitat being used by each bird observed will be recorded whenever possible. These data on the habitats being used by landbirds and shorebirds at the time of observation also will be used as an additional ground-reference data set to help in the mapping of wildlife habitats in the upper and middle Susitna basin (see Section 11.5).

As noted above, the point-count plot locations in 2013 will be selected using a pseudo-stratified random plot allocation procedure based on aerial photosignatures as the sampling strata, because it is unlikely that a current and complete habitat map will be available by spring 2013. In 2014, point-count locations will be selected again using a pseudo-stratified random plot allocation procedure, but mapped wildlife habitat types are expected to be available for use as the sampling strata (to the extent the wildlife habitat mapping is complete by spring 2014). In both years, the plot allocation will be constrained so that an adequate number of plots are placed in each mapped habitat or photosignature type. Without this constraint, an excessive number of plots would be located in the most common habitat types and far fewer would occur in uncommon types, resulting in under-sampling of uncommon habitat types. In all cases, sample points will be located in a random and spatially unbiased fashion (using the Geographic Information System [GIS]) within each mapped habitat or photosignature type, subject to the restriction of maintaining a minimum distance of 500 meters (1,640 feet) between sample points in open habitats and 250 meters (820 feet) in closed habitats. This sampling scheme will result in a selection of point-count locations that is unbiased with respect to the distribution of breeding birds on the landscape. The goal in plot allocation is to derive a set of sample points that are spread broadly across the study area and are replicated within each photosignature/habitat type to capture spatial variability in habitat use by breeding birds. Replicate sampling is also important to locate the often patchy occurrences of the less common species of conservation concern.

The point-count observers will estimate distance to birds detected on the surveys, using distance classes of 10-meter increments to 100 meters, then 100–125 meters, 125–150 meters, and >150 meters. Laser rangefinders will be used to confirm and calibrate the distance estimates. Using finer-scale distance classes in the field will allow the data to be grouped into broader categories later for analysis, if necessary. Experienced point-count observers will be used for these surveys, and all observers will receive refresher training before beginning fieldwork, to include bird identification (visual and by song and call), distance estimation to known targets, distance testing in a simulated survey, accuracy re-testing of distance estimates, and final distance testing in a simulated survey.

One extended field survey is planned in each summer breeding season in 2013 and 2014. In general, the survey is planned to begin in mid-May in each year, although survey timing may need to be adjusted to account for variability in the onset of breeding activities in response to early or late snowmelt and/or unseasonable temperature conditions in spring; local weather conditions and the activities of breeding birds in the Project area will be monitored each spring by communicating with the Project helicopter pilots and other wildlife researchers in the area (e.g., raptor biologists who will be surveying in the area in late April of each year). The surveys in May will be focused on breeding shorebirds and early nesting landbirds such as the Rusty Blackbird, a species of conservation concern in Alaska (USFWS 2008). It is likely that data on

early nesting resident birds also will be collected in the early portion of the survey period because nesting starts later at the higher elevations typical of the Project area. The point-count survey period will extend into mid-June to focus on neotropical migrant landbirds. Late-arriving flycatchers (e.g., Alder Flycatcher) will be present by early June. It is expected that some data on nesting resident birds and shorebirds will be collected during early June as well.

For the early portion of the survey period in May, point-count plots will be allocated preferentially in open habitats that are used by breeding shorebirds. These habitats include open wetlands in forested areas as well as open, dwarf scrub-dominated habitats in upland and alpine terrain. Woodland bog and tall scrub habitats in poorly drained lowlands will also be sampled because they are used by breeding shorebirds and Rusty Blackbirds. By later in the survey period (late May and early June), point-count plots will be allocated across all available habitats in the study area. As noted above, this survey will focus on neotropical migrant landbirds.

#### *10.16.4.1.3 Data Analysis*

Point-count survey data are subject to errors resulting from species-specific variability in vocalizations and behavior, observer variability in detecting birds, variability in detecting birds in different habitats, and variability in distance estimates. Consequently, to improve estimates of abundance and density, in analyses of the point-count survey data, researchers will use removal sampling (to improve the estimates of detection probabilities, especially for uncommon species; Farnsworth et al. 2002) and distance sampling (to calculate detection probabilities and densities; Buckland et al. 2001; Rosenstock et al. 2002), as recommended by USFWS during the landbird-shorebird meeting on September 6, 2012 (see Appendices 3 and 4). Observations will be stratified temporally (by point-count period) and spatially (by distance category) to estimate detection probabilities for all species, and allow corrections of the data to account for those birds that were present but not detected (Buckland et al. 2001; Rosenstock et al. 2002). Conducting both removal and distance analyses to estimate detection probabilities will result in improved estimates of abundance and density. As recommended by USFWS (see Appendices 3 and 4), existing detection functions, developed in other point-count studies in Alaska, will be used when necessary (e.g., to improve detectability estimates of uncommon species for which few observations in the study area may be available for analysis).

#### *10.16.4.2. Riparian- and Lacustrine-focused Surveys*

In May, an additional set of point-count surveys and linear surveys between point-count plot locations will be conducted specifically in riparian and lacustrine habitats that are expected to be affected by Project development. These surveys will be conducted in the inundation zone, in the infrastructure area surrounding the site of the proposed dam, and immediately below the location of the proposed dam. The surveys will target species (e.g., Semipalmated Plover, Spotted Sandpiper, Solitary Sandpiper, Wandering Tattler, Belted Kingfisher, and American Dipper) that are known to use riparian and lacustrine habitats, and which are not often recorded on point-counts allocated randomly across all available habitats. In addition to the point-count surveys in riparian and lacustrine habitats, researchers will walk along the sampled stream courses and lake/pond shorelines as they move between point-count locations. During these linear surveys, all birds observed will be recorded. The resulting data will be represented as the number of observations of each species per unit time spent in-transit to provide a standardized, relative measure of abundance across all species (following methods used by Andres et al. 1999 and

Boisvert and Schick 2007). A similar survey design for recording landbirds and shorebirds and relative levels of abundance while in transit between point-count locations was successfully completed on the Seward Peninsula in 2006 (Boisvert and Schick 2007) by the same wildlife contractors who will conduct the riparian- and lacustrine-focused surveys for this Project.

An additional goal of the riparian- and lacustrine-focused surveys will be to collect data on the distribution and abundance of piscivorous species (primarily Belted Kingfisher, but also American Dipper and Spotted Sandpiper, which occasionally consume fish) in the inundation zone and immediately below the location of the proposed dam (because elevated mercury levels could occur there in any fish that make it through the turbines). This information will be provided to the researchers conducting the mercury assessment study (see Section 5.7).

#### *10.16.4.3. Survey of Colonially Nesting Swallows*

The focal species for this survey effort include Bank Swallow, Cliff Swallow, and Violet-green Swallow. These three species are gregarious, colonial nesters that prefer riparian cutbanks and cliffs near lacustrine or fluvial waters (Kessel et al. 1982; Brown et al. 1992; Brown and Brown 2002; Garrison 2002). Bank and Violet-green swallows nest in burrows in relatively soft, sandy substrates and may form mixed-species colonies (Brown et al. 1992; Garrison 2002); data on Belted Kingfishers also are likely to be obtained incidentally because they nest in burrows in the same type of bank habitat. Cliff Swallows build nest cavities of mud and clay on rocky cliffs, bridges, and other human-made structures (Brown and Brown 2002). All three species feed on flying insects and often forage over or near water bodies (Brown et al. 1992; Brown and Brown 2002; Garrison 2002).

The swallow survey will be conducted using a motorized river boat for access to the Susitna River and tributary streams in the reservoir inundation zone. Researchers will search suitable nesting habitat and record nesting colonies. Two observers, plus the boat operator, will conduct the survey. In portions of tributary streams not accessible by boat, the survey will be conducted on foot. One observer will record the geographic locations of nesting habitat and colonies on a topographic map or aerial imagery, as well as in a hand-held global positioning system (GPS) receiver. The other observer will photograph the habitat and colonies to aid in accurate nest counts and to quantify the total area of potential nesting habitat. Both observers will assist in species identification and will estimate abundance and activity. Researchers will stop and observe active colonies for a minimum of 15 minutes, as recommended by Garrison et al. (1989), to allow accurate species identification and to obtain estimates of abundance and activity.

Accessible colonies will be closely inspected to estimate the proportion of active burrows. Researchers will check a stratified-random (spatially balanced) sample of burrows within accessible colonies using a flashlight and an angled mirror attached to an extendable rod. Burrows containing eggs, young, or adults will be considered occupied. The number of eggs or young in nests will be recorded. These data will be used to estimate the total number of active burrows in each colony and in the surveyed area as whole, thereby facilitating an estimation of population size.

The survey will be conducted in late June or early July of 2013 and 2014, depending on the snowmelt and temperature patterns in spring and early summer of each year. All three focal species normally arrive in Interior Alaska by mid- to late May and nesting occurs in June and July (Kessel et al. 1982; Brown et al. 1992; Brown and Brown 2002; Garrison 2002). Bank

Swallows may reuse old burrows or create new burrows during courtship (Hickman 1979). Therefore, surveys will be timed to occur after mating and nest establishment to reduce variability in burrow numbers (Jones 1987). Vacant, but suitable, nesting habitat will be recorded to quantify the total area of potential nesting habitat present in the study area and to inform subsequent survey efforts in 2014, which will focus on revisiting colonies located in 2013.

All accessible navigable portions of the study area will be surveyed once in 2013 and again in 2014. Changes in numbers of individuals, colonies, and nests between the two years could provide information on the variability in swallow populations and will improve abundance estimates. In addition to swallows, researchers will record incidental observations of other birds during this survey to support this and the other avian studies.

#### *10.16.4.4. Migration Surveys*

The migration survey component of this study will employ a combination of daytime visual sampling and nocturnal radar and visual sampling (using night-vision devices) during both the spring (late April to June) and fall (late August to mid-October) migration periods. This study component will be conducted in concert with the waterbirds study and is described in more detail in that study plan (see Section 10.15).

#### *10.16.4.5. Integration of Existing Information with Current Study*

The landbird and shorebird data collected in the APA Susitna Hydroelectric Project area in the 1980s (Kessel et al. 1982; AEA 2011) will be reviewed and incorporated into analyses of habitat use by these species presented in the Initial Study and Updated Study reports (see below). The primary focus will be to compare habitat-use patterns in the historical data with the results of current data analyses. The abundance and distribution information for landbirds and shorebirds from Kessel et al. (1982) will be reviewed to evaluate changes in abundance and distribution over the intervening 30 years. These historical comparisons will provide information on the recent trends for these species in the Project area, which will be useful for impact predictions and assessments.

#### *10.16.4.6. Mercury Assessment*

Scientific literature on the foraging habits and diets of piscivorous landbirds and shorebirds (primarily Belted Kingfisher, but also American Dipper and Spotted Sandpiper) will be reviewed to inform the mercury risk-assessment work (see Section 5.7, Mercury Assessment and Potential for Bioaccumulation Study) and to complement the field data gathered on the distribution and abundance of these species in the study area. To the extent possible, the information gathered will be focused on data from Alaska studies. In addition to the literature review, feathers will be collected from any kingfisher nests located during the swallow survey and will be provided to the study lead for the mercury study for laboratory analysis of methylmercury levels.

#### *10.16.4.7. Reporting and Data Deliverables*

The database and reporting deliverables for this study include the following:

- **Electronic copies of field data.** A geospatially-referenced relational database will be prepared, containing historical (APA Susitna Hydroelectric Project) data and current data



collected during this study, including representative photographs of breeding bird habitats at point-count locations. Naming conventions of files and data fields, spatial resolution, map projections, and metadata descriptions will meet the data standards established for the Project.

- **Initial Study Report and Updated Study Report.** The landbird and shorebird study results will be presented in the Initial and Updated Study reports, according to the schedule indicated in Table 10.16-1. The reports will include descriptions of the field survey methods, a map of the locations surveyed, and survey results with tables indicating abundance by species and habitat type.

#### **10.16.5. Consistency with Generally Accepted Scientific Practice**

The landbird and shorebird study will employ point-count surveys and analytical methods that conform to currently accepted, standard protocols for the monitoring of landbirds in remote, roadless landscapes of Alaska (Handel and Cady 2004; ALMS 2010). In recent years, these methods have also been adopted for shorebird surveys in Alaska (ASG 2008) and are especially appropriate in forested landscapes, where shorebirds typically occur in low densities and where plot-based methods would yield few observations, even with a relatively large survey effort.

The survey of colonially nesting swallows will employ the best available techniques. There currently is no standard protocol for surveying swallow colonies, but the methods proposed above follow those used elsewhere (Garrison et al. 1989; Watts et al. 1996). These methods are intended to be an improvement on point-count surveys, which are widely regarded to be inadequate for swallows and similar riparian species (Swanson and Nigro 2003).

The migration surveys will employ well-developed techniques for radar and visual sampling that have been used for similar studies in Alaska and elsewhere in North America, as is described in more detail in the waterbirds study plan (Section 10.15).

#### **10.16.6. Schedule**

The landbird and shorebird study will be a two-year effort. The schedule for the 2013 and 2014 activities is presented graphically below (Table 10.16-1). Logistics and survey planning will occur in March and April of 2013 and 2014. The radar and visual surveys of spring migrant birds (conducted in conjunction with the waterbirds study) will occur from late April to early June of 2013 and 2014. Point-count surveys and the riparian- and lacustrine-focused surveys will take place over a continuous 30-day period from mid-May to mid-June in 2013 and 2014. The swallow colony survey will take place over approximately four days between June 20 and July 10 of 2013 and 2014. Fall migration sampling (radar and visual sampling conducted in conjunction with the waterbirds study) will occur from mid-August to mid-October of 2013 and 2014. Data analysis will occur from July to December of 2013 and 2014. The Initial Study Report will be completed in February of 2014 and the Updated Study Report in February of 2015.

In 2014 and 2015, licensing participants will have opportunities to review and comment on the study reports (Initial Study Report in early 2014 and Updated Study Report in early 2015). Updates on the study progress will be provided during Technical Workgroup meetings, which will be held quarterly in 2013 and 2014.

### **10.16.7. Relationship with Other Studies**

This study has limited information needs from other studies, as is portrayed below (Figure 10.16-2). Before the point-count surveys and the riparian- and lacustrine-focused surveys, preliminary interpretation of aerial imagery from the Vegetation and Wildlife Habitat Mapping Study in the Upper and Middle Susitna Basin (Section 11.5) will be conducted to establish a preliminary classification of habitats for sample allocation. A pseudo-stratified random sampling scheme will be used to allocate point-count locations and effort among habitats. Habitat-specific survey data from the field surveys will be used to describe the distribution and abundance of each species detected in each habitat sampled, and density estimates will be calculated for each species in each habitat in which it was detected. Field survey data from the swallow colony survey will be used to document the locations of swallow colonies, which will facilitate separate population estimates of those species within the surveyed area. The diurnal visual and nocturnal radar migration surveys will be conducted in conjunction with the waterbird study (Section 10.15) to provide data on landbird and shorebird migration, including timing, volume, altitude, flight directions, and any corridors identified. High-value habitats for each species will be identified and density and distribution data from the various surveys of landbirds and shorebirds will be used to inform the Evaluation of Wildlife Habitat Use Study (Section 10.19), which in turn will be used to quantify potential Project impacts and to develop any protection, mitigation, and enhancement (PM&E) measures for landbirds and shorebirds, as appropriate (see below).

Landbirds and shorebirds could be affected directly by the loss of breeding habitat from the placement of fill and from the conversion of terrestrial habitats to lacustrine habitats in the proposed reservoir. Additional indirect impacts could occur from alteration of habitats by erosion, fugitive dust accumulation, permafrost degradation, landslides, and off-road vehicle use. Disturbance effects (displacement from breeding habitats) from construction and operations activities represent another possible source of indirect impacts. Direct impacts could occur through injury and mortality in various ways (e.g., if exposed to fuel from accidental spills or from in-flight collisions with infrastructure). Alterations in riparian wildlife habitats downstream from the proposed dam due to changes in instream flow, ice processes, and riverine geomorphology in the Susitna River are also possible, and will be addressed in the Riparian Vegetation Study Downstream of the Proposed Susitna-Watana Dam (see Section 11.6).

The impact assessment for landbirds and shorebirds will be conducted during preparation of AEA's License Application in 2015 by first conducting habitat-use evaluations (see Sections 10.19 and 11.5) to assign habitat values for each landbird and shorebird species to each of the wildlife habitats mapped in the Vegetation and Wildlife Habitat Mapping Study (see Section 11.5). Then the various components of the Project "footprint" will be overlaid on the mapped wildlife habitat types using GIS to quantify the acreages of important breeding habitats for each species that would be lost directly to fill or inundation. The determination of acreages of landbird and shorebird habitats that may be affected indirectly by habitat alteration and behavioral disturbance will be conducted similarly by overlaying habitat alteration and disturbance buffers (surrounding the proposed Project infrastructure). The size and number of habitat alteration and disturbance buffer(s) to be used will be determined based upon the final specifications for Project construction, operations, and maintenance activities, which will be provided in the Project description. Direct impacts on landbirds and shorebirds will be assessed qualitatively by evaluating the likelihood of injury and mortality from various sources during Project construction and operations. Cumulative effects on landbirds and shorebirds in the region of the

proposed Project will be assessed by evaluating the extent of the direct and indirect impacts expected from the Project in conjunction with the existing impacts on landbirds and shorebirds in the region.

#### **10.16.8. Level of Effort and Cost**

The ground-based point-count surveys are planned to be conducted over two years (2013–2014). The point-count field surveys (late spring and early summer) will be conducted each year by a crew of eight observers (four crews of two persons each). Point-count surveys will be conducted for approximately 28 days each year, with the goal of obtaining at least 800 point-count samples each year. Helicopter support will be required for this survey with drop-off and pick-ups each day in the field. The surveys will start at first light in the morning, which in the Project area will be approximately 3:30 A.M. The bulk of the costs associated with this study are for field sampling, data analysis, and reporting.

The survey of colonially nesting swallows will focus on estimating the abundance and distribution of swallows in riparian habitats within the inundation zone. The field survey will be conducted in a short time period (estimated four days each year) and require only two observers (plus a boat operator). However, this survey has specific field equipment and safety requirements, including boating and camping equipment that will require helicopter sling transport.

The estimated cost for the landbird/shorebird component of the spring and fall migration surveys is included in the cost estimate for the waterbirds study plan (see Section 10.15), so is not listed here.

The projected annual cost of this study is in the range of \$425,000–\$450,000, for an estimated total of \$850,000–\$900,000 over both years.

#### **10.16.9. Literature Cited**

- AEA (Alaska Energy Authority). 2011. Pre-Application Document: Susitna–Watana Hydroelectric Project FERC Project No. 14241. December 2011. Prepared for the Federal Energy Regulatory Commission by the Alaska Energy Authority, Anchorage.
- ALMS (Alaska Landbird Monitoring Survey). 2010. Boreal Partners in Flight: Alaska Landbird Monitoring Survey. Available online: <http://alaska.usgs.gov/science/biology/bpif/monitor/alms.php#information> (accessed March 10, 2012).
- Andres, B. A., D. L. Brann, and B. T. Browne. 1999. Legacy Resource Management Program: Inventory of breeding birds on local training areas of the Alaska Army National Guard. Final report prepared for Alaska Army National Guard, Fort Richardson, AK, by U.S. Fish and Wildlife Service, Anchorage, AK. 104 pp.
- ASG (Alaska Shorebird Group). 2008. Alaska Shorebird Conservation Plan. Version II. Anchorage.
- Boisvert, J. H., and C. T. Schick. 2007. Breeding bird surveys, Stewart River Training Area, Alaska, 2006. Final report prepared for Alaska Army National Guard, Fort Richardson, AK, by ABR, Inc.—Environmental Research & Services, Anchorage, AK. 25 pp.

- Brown, C. R., A. M. Knott, and E. J. Damrose. 1992. Violet-green Swallow. No. 14 in A. F. Poole, P. Stettenheim, and F. B. Gill, eds. *The Birds of North America*. The Birds of North America, Inc., Philadelphia, PA.
- Brown, C. R., and M. B. Brown. 2002. Cliff Swallow. No. 149 in A. F. Poole, P. Stettenheim, and F. B. Gill, eds. *The Birds of North America*. The Birds of North America, Inc., Philadelphia, PA.
- Buckland, S. T., D. R. Anderson, K. T. Burnham, J. L. Laake, D. L. Borchers, and J. Thomas. 2001. *Introduction to Distance Sampling: Estimating Abundance of Biological Populations*. Oxford University Press.
- Farnsworth, G. L., K. H. Pollock, J. D. Nichols, T. R. Simons, J. E. Hines, and J. R. Sauer. 2002. A removal model for estimating detection probabilities from point-count surveys. *Auk* 119: 414–425.
- Garrison, B. A. 2002. Bank Swallow. No. 414 in A. F. Poole, P. Stettenheim, and F. B. Gill, eds. *The Birds of North America*. The Birds of North America, Inc., Philadelphia, PA.
- Garrison, B. A., R. W. Schlorff, J. M. Humphrey, S. A. Laymon, and F. J. Michny. 1989. Population trends and management of the Bank Swallow (*Riparia riparia*) on the Sacramento River, California. U.S. Department of Agriculture, Forest Service General Technical Report PSW-110.
- Handel, C. M., and M. N. Cady. 2004. Alaska Landbird Monitoring Survey: protocol for setting up and conducting point count surveys. U.S. Geological Survey, Alaska Science Center, Anchorage. Available online: [http://alaska.usgs.gov/science/biology/bpif/monitor/alms/ALMSprotocol\\_2004.pdf](http://alaska.usgs.gov/science/biology/bpif/monitor/alms/ALMSprotocol_2004.pdf) (accessed August 11, 2011).
- Hickman, G. R. 1979. Nesting ecology of Bank Swallows in interior Alaska. M.S. thesis, University of Alaska, Fairbanks. 78 pp.
- Jones, G. 1987. Colonization patterns in Sand Martins *Riparia riparia*. *Bird Study* 34: 20–25.
- Kessel, B. 1979. Avian habitat classification for Alaska. *Murrelet* 60: 86–94.
- Kessel, B., S. O. MacDonald, D. D. Gibson, B. A. Cooper, and B. A. Anderson. 1982. Susitna Hydroelectric Project environmental studies, Phase I final report—Subtask 7.11: Birds and non-game mammals. Report prepared by University of Alaska Museum, Fairbanks, and Terrestrial Environmental Specialists, Inc., Phoenix, NY for Alaska Power Authority, Anchorage. 149 pp.
- Ralph, C. J., S. Droege, and J. R. Sauer. 1995. Managing and monitoring birds using point-counts: standards and applications. Pp. 161–168 in C. J. Ralph, S. Droege, and J. R. Sauer, eds. *Monitoring bird populations by point-counts*. U.S. Department of Agriculture, Forest Service General Technical Report PSW-GTR-149.
- Rosenstock, S. S., D. R. Anderson, K. M. Giesen, T. Leukering, and M. F. Carter. 2002. Landbird counting techniques: current practices and an alternative. *Auk* 119: 246–253.
- Swanson, S. A., and D. A. Nigro. 2003. A breeding landbird inventory of the Yukon-Charley Rivers National Preserve, Alaska, June 1999 and 2000. In-house Report YUCH-03-01. National Park Service, Fairbanks, Alaska.

- USFWS. 2008. Birds of Conservation Concern. 2008. Division of Migratory Bird Management, Arlington, VA. 85 pp. Available online: <http://www.fws.gov/migratorybirds> (accessed July 12, 2011).
- Watts, B. D., M. A. Byrd, and M. U. Watts. 1996. Status and distribution of Cliff Swallows in coastal Virginia. *The Raven* 67: 21–24.

**10.16.10. Tables**

**Table 10.16-1. Schedule for implementation of the landbird and shorebird study.**

Activity	2013				2014				2015
	1 Q	2 Q	3 Q	4 Q	1 Q	2 Q	3 Q	4 Q	1Q
Review aerial imagery & select point-count survey sites	—				—				
Point-count field surveys		—				—			
Riparian- and lacustrine-focused surveys		—				—			
Swallow colony survey			—				—		
Migration surveys (radar & visual)		—	—			—	—		
Data analysis			—	—			—	—	
Initial Study Report					△				
Updated Study Report									▲

Legend:

- Planned Activity
- △ Initial Study Report
- ▲ Updated Study Report

10.16.11. Figures




Figure 10.16-1. Landbird and shorebird study area.

**STUDY INTERDEPENDENCIES FOR LANDBIRD AND SHOREBIRD STUDY**




Figure 10.16-2. Study interdependencies for the landbird and shorebird study.



## **10.17. Population Ecology of Willow Ptarmigan in Game Management Unit 13**

### **10.17.1. General Description of the Proposed Study**

The Ptarmigan Study, which will be conducted by the Alaska Department of Fish and Game (ADF&G), will be a two-year investigation, beginning in spring 2013 and continuing through winter 2014 to inform the FERC licensing process. To take full advantage of the expected life span of the radio telemetry tags that will be deployed for the study, ADF&G may continue to track tagged birds further (into 2016), but the data obtained in 2013 and 2014 will be sufficient for the FERC licensing effort.

#### **Study Goals and Objectives**

The goal of this study is to provide the necessary data to evaluate the potential effects of the proposed Project on Willow Ptarmigan, the predominant species of upland game bird in the Project area and surrounding areas. The area of interest consists of Subunits 13A and 13E of Game Management Unit (GMU) 13 (Figure 10.17-1).

The study has four objectives:

- Determine the seasonal distribution of Willow Ptarmigan in the Project area.
- Determine the seasonal migratory patterns of Willow Ptarmigan that occur in the Project area.
- Estimate the abundance of ptarmigan in the Project area during the breeding season and during the fall.
- Estimate seasonal survival of Willow Ptarmigan.

The data gathered during the study will be integrated to determine potential effects of the Project on Willow Ptarmigan.

### **10.17.2. Existing Information and Need for Additional Information**

The Willow Ptarmigan is the most common and widespread ptarmigan in Alaska, constituting an estimated 65–70 percent of all ptarmigan statewide, followed by Rock Ptarmigan at 25–30 percent, and White-tailed Ptarmigan at <10 percent (Taylor 1994). All three ptarmigan species occur in GMU 13 (Taylor 2000).

Ptarmigan hunting is a very popular activity in the fall and winter months in GMU 13 due to the accessibility of the unit from the state highway system. Since 1997, ADF&G has conducted ptarmigan surveys in spring along the Denali, Parks, and Richardson highways to quantify the relative abundance of territorial males. All survey efforts have centered on road-accessible areas within GMU 13. Those surveys suggest that Willow Ptarmigan along the road system portions of GMU 13 are declining in abundance or have remained at low abundance since 2000 (Bill Taylor, pers. comm.). Due to this continued low abundance, ADF&G recommended that the Alaska Board of Game reduce the bag limit of ptarmigan from 10 per day to 5 per day in Subunits 13A, 13B, and 13E between December 1 and March 31, and this recommendation took effect during the 2005–2006 regulatory year. Continued low abundance resulted in further harvest restrictions in Subunit 13B, and beginning in 2009, the ptarmigan season has been closed after November 30

each year. ADF&G has been unable to commit additional resources to better understand the life history of GMU 13 ptarmigan populations and there is little information on the habitat value of the Project area for ptarmigan.

Ptarmigan that winter in the Project area may be either resident or migratory birds. To better predict the potential effects of the proposed Project on Willow Ptarmigan, information needs to be collected to determine the annual ranges of ptarmigan that may use habitats in GMU 13. In particular, information is needed to evaluate the relative importance of the Project area to resident and migratory ptarmigan and the seasonal ranges of migratory birds need to be delineated.

### **10.17.3. Study Area**

Willow Ptarmigan will be captured within a 15-mile buffer around the proposed dam site and reservoir and the access and transmission corridor alternatives (Figure 10.17-1). Capture locations will be in the headwaters of several major river drainages. The study area is composed of alpine habitats at higher elevations and subalpine spruce habitats at lower elevations. Areas in which Willow Ptarmigan will be captured are roadless, although periodic, but infrequent, all-terrain vehicle use can occur year-round.

The areas selected for capture have been identified previously as locations with relatively high breeding densities of Willow Ptarmigan. Initial capture efforts will focus on three areas, including upper Fog Creek (tributary to the upper Susitna River), upper Busch Creek (tributary to Goose Creek), and the pass between upper Jay and Coal creeks.

Radio-tagged Willow Ptarmigan are expected to remain within 50 miles of the original capture site, although movements may exceed that distance in some cases (Irving et al. 1967). Aerial surveys to locate birds with radios will be conducted in appropriate habitats within 50 miles of the original capture locations.

### **10.17.4. Study Methods**

#### *10.17.4.1. Capture and Radio-tagging of Ptarmigan*

Beginning in April 2013, 50–100 Willow Ptarmigan will be captured annually at three sites within Subunits 13A and 13E (Figure 10.17-1) and fitted with radio-transmitter necklaces. All three sites are within 15 miles of either the proposed reservoir inundation zone or the access corridors. Alternative capture sites may be needed based on conditions each spring, depending on factors such as ptarmigan abundance, snow depth, and fixed-wing airplane access. Potential alternative capture sites (Figure 10.17-1) will be considered during field operations in the spring and summer of 2013.

Capture sites and future alternative sites have and will be identified based on several criteria:

- Willow Ptarmigan abundance
- Proximity to the future reservoir or access routes
- Ease of access using either fixed-wing or helicopter
- Observed springtime conditions (i.e., snow depth, and habitat availability during the capture time period)

During the breeding season in April and May of each year, several two-person teams will be deployed to various capture locations using wheel-ski equipped fixed-wing aircraft. Teams will attempt to capture 25–50 Willow Ptarmigan during the spring effort. Several teams will again be deployed in mid-August to September, using fixed-wing aircraft, in an effort to capture an additional 25–50 adult and fledged juveniles before brood dispersal occurs (Weeden and Watson 1967).

When capture efforts begin in April, male Willow Ptarmigan will be located visually or by using a playback recording of a territorial male Willow Ptarmigan (Taylor 1999; Peyton 1999; Savage et al. 2011). Playback recordings will be used effectively under low wind conditions ( $\leq 5$  miles per hour) with no precipitation during early morning or late evening hours. Once ptarmigan are known to be in the vicinity, a styrofoam decoy and remotely powered caller will be placed within the defensive range ( $<100$  meters [328 feet]) of a territorial male. A mist net will be deployed around the decoy and remotely powered caller in an attempt to capture the territorial male when he responds defensively to the call and decoy. Mist nets designed by Avinet ([www.avinet.com](http://www.avinet.com)) for capturing small hawks and large shorebirds will be used (Silvy and Robel 1968). These black nylon nets have a 100-millimeter (~4-inch) mesh and are 2.6 meters (8.5 feet) tall, with four shelves (Browsers and Connelly 1986). When circumstances allow during spring capture efforts, a hand-held Coda net gun ([www.codaenterprises.com](http://www.codaenterprises.com)) with a 12-square-foot net and 3-inch mesh will also be used opportunistically to capture territorial male birds on the ground, primarily as a backup to the mist net method. This method has been used from a helicopter to capture short-eared owls in northern Alaska and has proven to be safe and effective (T. Booms, ADF&G, pers. comm.). The use of decoys and calls is a novel adaptation to attempt to increase the number of captures typical of previous netting methods ( $>30$  ptarmigan annually; Skinner et al. 1998; Kaler et al. 2010). No attempt to capture nesting or brood-rearing females will occur.

Post-breeding resident and migrant birds will be targeted for capture during a second annual capture effort in mid-August through September. Flocks of ptarmigan will be located visually, mist nets will be strategically placed around or in the vicinity, and ptarmigan of all age/sex classes will be flushed into the mist nets. Fall captures will be similarly outfitted with radio-transmitter necklaces.

At least two people will be present for any single capture event to remove birds from mist nets, handle, and release birds as quickly as possible. After capture, Willow Ptarmigan will be restrained in a capture bag or by holding their wings against their bodies. Birds will be instrumented with a necklace-mounted A3950 VHF radio transmitter with a 10-inch whip antenna (Raymond 1999; Paragi et al. 2012; Figure 10.17-2) from Advanced Telemetry Systems (ATS, [www.atstrack.com](http://www.atstrack.com)). The entire radio and necklace package will weigh up to 10.7 grams (0.4 ounce) (1.7 percent of the body mass based on known weights of hunter-harvested Willow Ptarmigan; Hudson 1986; Thirgood et al. 1995). Radios will transmit in the frequency range of 148.000 Mhz. The transmitter is secured by a rubber-sheathed wire fitted over the bird's neck and crimped on either end to ensure its fit (Figure 10.17-2). The transmitter will be adjusted to compensate for crop expansion. No tissue samples will be collected from captured Willow Ptarmigan. Birds will be handled for 5–10 minutes and released at their point of capture.

Age and sex, based on plumage characteristics (Bergerud et al. 1963; Weeden and Watson 1967; Braun and Rogers 1971; Hudson 1986) will be recorded for each bird captured. Individually numbered leg bands will be placed on each radio-tagged bird. These tags will be useful for ground observations and to identify human-harvested birds or prey remains that may be found

during field efforts. A Global Positioning System (GPS) receiver will be used to record the location of capture. Date, time, and weather conditions also will be recorded. If a territorial male is captured, an attempt will be made to identify and record the location(s) of his territory post(s).

Radio tags will not be removed at the conclusion of the study, nor will tags drop off. There is little evidence to suggest that radio tags have a negative effect on the survival or breeding success of ptarmigan and other galliformes (Thirgood et al. 1995; Palmer and Wellendorf 2007; Terhune et al. 2007). Radio-tagged Willow Ptarmigan will be closely monitored within 24 hours of capture to document capture myopathy or other obvious handling-induced stresses. All potential capture and marking methods will be fully evaluated and compliant with Alaska Interagency Animal Care and Use Committee (IACUC) certification. ADF&G will ensure compliance with all IACUC policies.

#### *10.17.4.2. Relocation of Radio-tagged Ptarmigan*

Radio-tagged ptarmigan will be relocated during aerial surveys conducted throughout the year to record habitat use, movements, and mortality. Birds will be tracked and relocated using a fixed-wing airplane equipped with wheel-skis, which will decrease search time and increase the area that can be covered. The first aerial survey will be performed within 10 days of capture to document survival rates of the birds recently radio-tagged. At least six additional aerial surveys will be performed annually: two in late summer (August–September), two in mid-winter (November–February), and two in early spring (late March to mid-April).

Range of radio tags will be tested before deployment. However, temperature may affect transmission range (T. Paragi and B. Taylor, ADF&G, pers. comm.). Therefore, to ensure a systematic search pattern, aerial surveys will be flown using a preselected 5-mile grid and flown at an altitude of 1,500–2,000 feet within Subunits 13A and 13E.

An ATS 4520 receiver will be used to locate radio-tagged birds. Two 4-element Yagi antennas will be mounted to each strut of the aircraft. A GPS receiver mounted at the windshield of the aircraft and connected to the ATS 4520 receiver will provide a location for each data record. Upon completion of each aerial survey, receivers will be downloaded to a field laptop or Local Area Network (LAN) at the ADF&G office in Palmer for future analysis and specific location determination of each tagged bird.

#### *10.17.4.3. Aerial Transect Surveys*

During September and March each year, aerial transect surveys will be flown to assess the abundance and density of ptarmigan using line-transect or repeat-count techniques (Royle and Dorazio 2008; Thomas et al. 2010). In addition to abundance, these surveys will provide data on the overall distribution of all ptarmigan (not just radio-tagged individuals) in Subunits 13A and 13E.

#### *10.17.4.4. Analysis of Radio Telemetry Data*

After the radio receivers have been downloaded, data will be transferred to a Microsoft Access database for analysis. Maps will be created using Geographic Information System (GIS) software (ArcMAP) for each aerial survey day, indicating the location of each relocated Willow Ptarmigan. These data will be catalogued and used for spatial analyses.

Movement and survival rates of tagged birds will be estimated using multistate models (Brownie et al. 1993). Occupancy models of aerial survey data will be used to estimate the probability that an area is used and to identify changes in the probability of use between fall and spring surveys (Nichols et al. 2008).

The combination of telemetry transmitters and large-scale aerial surveys will provide both specific information on individual movements and habitat use and general information on species distribution. These survey techniques are being developed and implemented for another study of ptarmigan north of the Brooks Range (K. Christie, pers. comm.).

#### **10.17.5. Consistency with Generally Accepted Scientific Practice**

Habitat availability and use analyses allow an ecosystem approach to impact assessment, and GIS-based analysis has become a standard and straightforward method of evaluating the impacts of habitat loss and alteration. Ptarmigan captures will be conducted by adapting fairly standard capture methods to the situation. With continuous improvements in technology, particularly in battery and transmitter weights, radio telemetry is an important and increasingly standard method of obtaining movement data even for small birds and mammals. All potential capture and marking methods will be fully evaluated and compliant with Alaska Interagency Animal Care and Use Committee (IACUC) certification. ADF&G will ensure compliance with all IACUC policies. There is little evidence to suggest that radio tags have a negative effect on the survival or breeding success of ptarmigan and other galliformes (Thirgood et al. 1995; Palmer and Wellendorf 2007; Terhune et al. 2007). Radio-tagged Willow Ptarmigan will be closely monitored within 24 hours of capture for signs of capture myopathy or other obvious handling-induced stresses.

#### **10.17.6. Schedule**

As is summarized in Table 10.17-1, aerial transect surveys will begin in March 2013 and ptarmigan tagging and tracking will begin in April 2013 and will continue through the end of 2014. ADF&G may continue to track tagged ptarmigan through 2016, corresponding to the anticipated lifespan of the radio transmitters. Project milestones will follow the same monthly schedule each year, unless noted otherwise. Aerial transect surveys will be flown in March and September. Capture of ptarmigan and deployment of radio tags will be conducted in April, May, and August, and radio-tracking will be conducted from August 2013 through May 2014 and from August through December 2014. Updates on the study progress will be provided during Technical Workgroup meetings which will be held quarterly in 2013 and 2014. The Initial Study Report will be completed by February 2014, and the Updated Study Report will be completed by February 2015.

#### **10.17.7. Relationship with Other Studies**

As is depicted in Figure 10.17-3, the ptarmigan study will not require information inputs from other Project studies. The efforts conducted for this study will consist primarily of capture, tagging, and tracking of radio-tagged birds and aerial transect surveys of distribution and abundance. Radio-tracking data will be used to create a geodatabase of locations (with sex and season attributes), which will be used to evaluate regional occupancy and to conduct multistate modeling of seasonal range use, movements, and survival. Information on movements will be

used to delineate the seasonal ranges of ptarmigan using the study area. Data from aerial transect surveys will provide additional details on the distribution and abundance of ptarmigan in the study area in late winter and fall. Location data from both radio-tracking and transect surveys will identify high-value habitats for ptarmigan in the study area, which will be used to inform the Evaluation of Wildlife Habitat Use (Section 10.19). The aggregate data obtained on abundance, density, seasonal distribution, and movements will be used to assess the potential impacts of the Project and to develop PM&E measures, as appropriate.

The Ptarmigan Study is designed to provide relevant information for assessing potential direct, indirect, and cumulative effects of the Project, which may include the following:

- Permanent habitat loss caused by Project facilities, including the reservoir, powerhouse, and other permanent Project facilities.
- Temporary loss or alteration of habitats affected by clearing, dust fallout, gravel spray, persistent snow drifts, impoundments, thermokarst, contaminant spills, and other indirect effects of Project construction and operation.
- Behavioral disturbance of ptarmigan by Project construction and operation activities, including vehicle and heavy equipment traffic, geophysical investigations, and other human activities in the Project area.
- Indirect habitat loss through displacement of birds that avoid Project facilities or transportation routes.
- Increased predation on birds or their eggs that may result from attraction of predators to anthropogenic foods or artificial structures (such as perches on power poles or power lines, for example).
- Injury and mortality of birds from collisions with aircraft, vehicles, or structures (such as power lines, for example).
- Injury and mortality of birds due to contact with or ingestion of contaminants (including fuels), including potential indirect effects on forage plants.
- Increased harvest of ptarmigan resulting from improvements in access to humans.

During the impact assessment that will be conducted for the FERC License Application in 2015, data on the distribution, abundance, movements, productivity, and habitat use of Willow Ptarmigan in the study area will be used to assess Project impacts through geospatial analysis and evaluation of the responses of the species to other similar projects, as documented in the scientific literature. Using GIS software, species abundance data recorded among different habitat types will be combined with the spatially explicit wildlife habitat map of the Project area that will be developed under the botanical resources study plans to assess direct and indirect impacts of habitat loss and alteration and behavioral disturbance. The direct and indirect impacts of the Project will be evaluated by overlaying the Project features (including the reservoir impoundment, related infrastructure areas, and access road and power transmission corridors), and the seasonal ranges of ptarmigan on the Project habitat map. Seasonal ranges will be delineated with radio telemetry, using the recorded movements of a sample of birds to which radios have been attached. By plotting ptarmigan locations on the habitat map, high-value or high-density habitats can be identified. Indirect impacts will be estimated by applying various buffer distances on Project features, as determined from the available information on the anticipated effects of construction disturbance and habitat-related changes due to infrastructure and development, and identifying areas of high-value habitats that are affected. The GIS analysis will be combined with results from the telemetry study and transect surveys, as well as from the

scientific literature, to estimate the geographic extent, frequency, duration, and magnitude of Project effects on ptarmigan. Any necessary protection, mitigation, and enhancement (PM&E) measures will be developed, as appropriate, by examining the distribution and abundance of Willow Ptarmigan among habitats in relation to the geographic extent and seasonal timing of Project activities.

#### **10.17.8. Level of Effort and Cost**

This study is a multi-year effort that will be conducted by ADF&G. The estimated cost of the study over both years is \$415,000.

#### **10.17.9. Literature Cited**

- Bergerud, A. T., S. S. Peters, and R. McGrath. 1963. Determining sex and age of Willow Ptarmigan in Newfoundland. *Journal of Wildlife Management* 27: 700–711.
- Braun, C. E., and G. E. Rogers. 1971. The White-tailed Ptarmigan in Colorado. Colorado Division of Game, Fish, and Parks, Technical Publication No. 27.
- Browsers, H. W., and J. W. Connelly. 1986. Capturing sage grouse with mist nets. *Prairie Naturalist* 18: 185–188.
- Brownie, C., J. E. Hines, J. D. Nichols, K. H. Pollock, and J. B. Hestbeck. 1993. Capture-recapture studies for multiple strata including non-Markovian transitions. *Biometrics* 49: 1173–1187.
- Hudson, P. J. 1986. *Red Grouse: The biology and management of a wild gamebird*. The Game Conservancy Trust, Fordingbridge, UK. 250pp.
- Irving, L., G. C. West, L. J. Peyton, and S. Paneak. 1967. Migration of Willow Ptarmigan in Arctic Alaska. *Arctic* 20: 77–85.
- Kaler, R. S. A., S. E. Ebbert, C. E. Braun, and B. K. Sandercock. 2010. Demography of a reintroduced population of Evermann's Rock Ptarmigan in the Aleutian Islands. *Wilson Journal of Ornithology* 122: 1–14.
- Nichols, J. D., L. L. Bailey, A. F. O'Connell, N. W. Talancy, E. H. C. Grant, A. T. Gilbert, E. M. Annand, T. P. Husband, and J. E. Hines. 2008. Multi-scale occupancy estimation and modeling using multiple detection methods. *Journal of Applied Ecology* 45: 1321–1329.
- Palmer, W. E., and S. D. Wellendorf. 2007. Effect of radiotransmitters on Northern Bobwhite annual survival. *Journal of Wildlife Management* 71: 1281–1287.
- Paragi, T. F., J. D. Mason, and S. M. Brainerd. 2012. Summer habitat selection by Sharp-tailed Grouse in eastern interior Alaska. Alaska Department of Fish and Game, Federal Aid in Wildlife Restoration, Final Research Report, Grants W-33-8 and W-33-9, Project 10.01, Juneau.
- Peyton L. J. 1999. *Bird songs of Alaska*. Library of Natural Sounds, Cornell Laboratory of Ornithology, Ithaca, NY.
- Raymond, R. L. 1999. Sharp-tailed Grouse habitat study in eastern Interior Alaska. Alaska Department of Fish and Game, Juneau.

- Royle, J. A., and R. M. Dorazio. 2008. *Hierarchical Modeling and Inference in Ecology: The Analysis of Data from Populations, Metapopulations, and Communities*. Academic Press, San Diego, CA. 444 pp.
- Savage, S. E., K. J. Payne, and R. T. Finer. 2011. Willow Ptarmigan pilot study, Alaska Peninsula, May 2011. Unpublished report, U.S. Fish and Wildlife Service. 40 pp.
- Silvy, N. J., and R. J. Robel. 1968. Mist nets and cannon nets compared for capturing prairie chickens on booming grounds. *Journal of Wildlife Management* 32: 175–178.
- Skinner, W. R., D. P. Snow, and N. F. Payne. 1998. A capture technique for juvenile Willow Ptarmigan. *Wildlife Society Bulletin* 26: 111–112.
- Taylor, W. P. 1994. Game Management Unit 13 ptarmigan hunter and harvest report, 1992–94. Unpublished report, Alaska Department of Fish and Game.
- Taylor, W. P. 1999. *Game Management Unit 13 ptarmigan population studies*. Alaska Department of Fish and Game. Juneau, AK.
- Taylor, W. P. 2000. *Game Management Unit 13 ptarmigan population studies*. Federal Aid in Wildlife Restoration, final research performance report, 1 August 1997–30 June 1999. Grants W-27-1 and W-27-2, Study 10.70. Alaska Department of Fish and Game, Juneau. 12 pp.
- Terhune, T. M., D. C. Sisson, J. B. Grand, and H. L. Stribling. 2007. Factors influencing survival of radiotagged and banded Northern Bobwhites in Georgia. *Journal of Wildlife Management* 71: 1288–1297.
- Thirgood, S. J., S. M. Redpath, P. J. Hudson, M. M. Hurley, and N. J. Aebischer. 1995. Effects of necklace radio transmitters on survival and breeding success of Red Grouse *Lagopus lagopus scoticus*. *Wildlife Biology* 1: 121–126.
- Thomas, L., S. T. Buckland, E. A. Rexstad, J. L. Laake, S. Strindberg, S. L. Hedley, J. R. B. Bishop, T. A. Marques, and K. P. Burnham. 2010. Distance software: design and analysis of distance-sampling surveys for estimating population size. *Journal of Applied Ecology* 47: 5–14.
- Weeden, R. B., and A. Watson. 1967. Determining the age of Rock Ptarmigan in Alaska and Scotland. *Journal of Wildlife Management* 31: 825–826.



**10.17.10. Tables**

**Table 10.17-1. Schedule for implementation of the Willow Ptarmigan Study.**

Activity	2013				2014				2015
	1 Q	2 Q	3 Q	4 Q	1 Q	2 Q	3 Q	4 Q	1 Q
First field season: capture and tag birds		—	—						
Aerial radio-tracking surveys			—	—	—	—	—	—	
Aerial transect surveys	—		—		—		—		
Initial Study Report				—	△				
Second field season: capture and tag birds						—	—		
Telemetry analyses (occupancy modeling, multistate models)					—	—	—		
Updated Study Report								—	▲

Legend:

- Planned Activity
- △ Initial Study Report
- ▲ Updated Study Report

10.17.11. Figures




Figure 10.17-1. Willow Ptarmigan study area.



**Figure 10.17-2. A Sharp-tailed Grouse equipped with an ATS 3950 radio tag identical to the model that will be used for Willow Ptarmigan. [Photo by Scott Brainerd, ADF&G, 2012.]**

### STUDY INTERDEPENDENCIES FOR WILLOW PTARMIGAN STUDY




Figure 10.17-3. Study interdependencies for the Willow Ptarmigan study.

## **10.18. Wood Frog Occupancy and Habitat Use**

### **10.18.1. General Description of the Proposed Study**

The Wood Frog Occupancy and Habitat Use study (Wood Frog Study) will be conducted over two years (2013 and 2014), with fieldwork scheduled in May each year. The study will focus on evaluating the distribution of breeding wood frogs in those portions of the Project area in the upper and middle Susitna basin where breeding frogs could be directly or indirectly affected by Project development activities. The study will involve both field surveys and habitat occupancy modeling. In addition, AEA is proposing to opportunistically capture and sample frogs (non-lethally) to assay for the presence of the chytrid fungus that has been linked to amphibian declines worldwide (see Section 10.18.2 below).

#### **Study Goal and Objectives**

The goal of the Wood Frog Study is to characterize the use of the Project area by breeding wood frogs to facilitate an assessment of potential impacts on wood frogs from development of the proposed Project.

The study has four objectives:

- Review existing data on habitat use and distribution of breeding wood frogs in a broad region surrounding the Project area.
- Estimate the current occupancy rate for breeding wood frogs in suitable habitats in the study area through a combination of field surveys and habitat-occupancy modeling.
- Use information on current habitat occupancy and habitat use to estimate the habitat loss and alteration expected to occur from development of the Project.
- Sample frogs opportunistically for the presence of the chytrid fungus that has been linked to amphibian population declines. (At the request of state and federal management agencies, EA has agreed to sample for the chytrid fungus, to opportunistically take advantage of planned fieldwork by providing some baseline information on the occurrence of the fungus in the Project area pre-development.)

The Wood Frog Study is planned as a two-year study. Results from the first year of work in 2013 will be presented in the Initial Study Report and will be used to update the study plan for 2014, as needed, and to adjust the field survey methods and survey areas, if necessary, based on comments on the Initial Study Report by FERC, resource agencies, and other licensing participants.

### **10.18.2. Existing Information and Need for Additional Information**

Because amphibians were not included in the original Alaska Power Authority (APA) Susitna Hydroelectric Project environmental program studies in the 1980s, data on the occurrence of wood frogs in the upper Susitna drainage is lacking. It is likely that wood frogs occur in the Project area because they occur in suitable habitats throughout southern Alaska and in the interior north to the southern slopes of the Brooks Range; they have also been documented in Denali National Park and Preserve, near Healy, and in the lower Susitna drainage (Cook and MacDonald 2003; Anderson 2004; Gotthardt 2004, 2005; Hokit and Brown 2006; MacDonald 2010). Amphibian populations appear to have been declining worldwide for several decades (Blaustein and Wake 1990; McCallum 2007) and, although populations may be healthy in

Alaska (Gotthardt 2004, 2005), concern has been expressed about the conservation status of wood frogs in Alaska (ADF&G 2006). Because of this and because their status in the Project area is unknown, field surveys for wood frogs will be conducted in areas likely to be affected by Project facilities and activities.

*Batrachochytrium dendrobatidis* (Bd) is a chytrid fungus that causes the disease chytridiomycosis in amphibians. Since it was first discovered in amphibians in 1998, it has devastated amphibian populations around the world, including in North America. Bd is sometimes a non-lethal parasite and some amphibian species and some populations of susceptible species are known to survive infection. The fungus is widespread and ranges from lowland forests to cold mountain tops, and is typically associated with host mortality in high altitude environments and during winter, with greater pathogenicity at lower temperatures. Wood frogs have been identified as a frog species susceptible to infection by Bd, and Bd was first detected in a dead wood frog in Kenai National Wildlife Refuge in 2002 (Reeves 2008). The only other positive detection of Bd was near Dyea in Southeast Alaska in 2006 and was associated with the apparent die-off of western toads in Southeast Alaska (Sunday, May 21, 2006, *Juneau Empire*). No sampling for Bd has occurred in the Project area. Bd is believed to spread mainly through contact between infected frogs or with infected water. In its comments on study requests for the Project, the Alaska Department of Fish and Game (ADF&G) requested that water or frogs at survey locations be tested for the presence of Bd (see Section 8.4 in AEA 2012).

### **10.18.3. Study Area**

The study area includes those water bodies and suitable wetland habitats in the proposed Project area in which habitat loss, habitat alteration, and disturbance are expected to occur. The proposed study area encompasses the reservoir impoundment zone, areas for infrastructure of the dam and powerhouse and supporting facilities, the proposed access roads, and material sites (Figure 10.18-1).

### **10.18.4. Study Methods**

#### *10.18.4.1. Field Surveys and Occupancy Modeling*

Because the study area is large and the calling period of breeding male frogs is short, this study will not involve a comprehensive survey of all potential frog breeding habitat present in the study area. Instead, observers will survey for frogs in smaller sampling regions containing suitable habitats. Up to 10 sampling regions will be selected to survey for wood frogs in the study area (two in each of the three access road corridors and four in the reservoir zone and dam and camp facilities area of the proposed reservoir). Using a Geographic Information System (GIS), the sampling regions will be selected randomly from available areas of suitable water body and wetland habitats for frogs by overlaying a grid onto the full study area (Figure 10.18-1), removing all grid cells that contain no suitable habitat (or very little habitat), and then randomly selecting from the remaining cells for sampling. The random cell-selection process will be stratified so that sampling regions are selected in each of the transmission line/access road corridors and in the inundation/dam infrastructure zone, as noted above. One exception to this random cell-selection process will be made to ensure that the water bodies and wetlands in

the dam and camp facilities area near the proposed dam are included as one of the sampling regions.

Within the study area boundary, potential water bodies and wetland habitats (with permanent standing water) to be surveyed will be identified from interpretation of aerial photos or remote-sensed imagery and from the preliminary mapping of vegetation, wildlife habitats, and wetlands (see Sections 11.5 and 11.7). From this set of water bodies and wetlands, habitats will be categorized as having a higher or lower probability of supporting breeding frogs. The random sampling of water bodies and wetlands in each sampling region then will be stratified so that a greater effort is made to survey habitats that have a higher probability of being used by frogs. Habitats more likely to be used by breeding frogs will be identified using GIS as those that (1) are not known to support fish (if available, data from the fish distribution and abundance studies [see Section 9] will be used to document fish occurrence, recognizing that those data may not be complete in 2013); (2) are not connected to stream systems supporting fish; and (3) have at least some emergent vegetation. Fish are efficient predators of frogs, and in studies in south-central Alaska, including portions of the lower Susitna basin (Gotthardt 2004), frogs have been recorded in lower numbers in water bodies that support fish. Emergent and aquatic vegetation in water bodies provides a substrate for frog egg-masses and escape cover from aquatic predators, as well as helping to increase dissolved oxygen in the water (France 1997; Babbitt and Tanner 1998).

With a set of water bodies and wetlands identified throughout the study area that have the potential to support breeding frogs, a subset of those sites in each of the 10 sampling regions in the study area will be selected to survey for breeding frogs. Within each sampling region, a minimum of 12 potential water body and/or wetland sites will be selected for sampling using a stratified random selection process in GIS, as noted above, so that more effort is expended on sampling sites likely to be of higher value to breeding frogs. With 12 sites sampled in each of 10 sampling regions, at least 120 sites are expected to be surveyed across the full study area in each year. In practice, more than 120 sites may be surveyed because some sites will not need to be visited twice (i.e., when frogs are detected on the first visit, see below). A minimum distance of 250 meters (820 feet) between sample sites will be maintained to avoid duplicate detections of frogs.

Ground-based auditory surveys of the randomly selected water bodies and wetlands in the study area will be conducted during the breeding season for frogs (mid- to late May). (In addition to these surveys, it is expected that incidental detections of wood frogs will also be documented during data collection efforts for other resources [e.g., fisheries, vegetation and wetlands, and ground-based bird surveys], and this information may provide additional information on the occurrence of frogs in the study area.) Survey sites will be accessed by helicopter and on foot by navigating to predetermined sample sites using hand-held global positioning system (GPS) receivers. The field surveys will involve auditory detections of calling frogs to ascertain the presence or absence of wood frogs at each sampling site. Observations along the margins of each water body or wetland will be made at locations where observers can readily hear calling frogs. For small water bodies and wetlands, a single observation point will suffice to detect the presence of frogs, but for large water bodies and wetlands, multiple observation points may be needed to determine the presence of frogs. For large water bodies and wetlands, up to four observation points will be located and sampled, with distances of up to 500 meters between each point to achieve adequate survey coverage. Up to two independent, replicate surveys will be made by trained observers to each water body during the peak calling period (approximately

1200 to 2200 hours) of male wood frogs in southern Alaska (Gotthardt 2004; PLP 2011). Due to variability in the calling frequency of male wood frogs even during the peak of the breeding season (see PLP 2011), two visits may be needed to detect frogs at some water bodies; these replicate survey data will also be used to calculate the detectability of calling frogs, which is a critical component of this study. The second survey at each site will be conducted by a different observer with no knowledge of the survey results from the first survey. However, because this study involves the use of a “removal design” to estimate occupancy, if detected on the first survey, a second survey will not be needed (i.e., that site will be “removed” from further sampling; see Mackenzie and Royle [2005] for more information). Surveys will be conducted only under favorable weather conditions (e.g., light rain or no rain, air temperature higher than 4° C [39° F], and wind speeds less than or equal to 25 kilometers per hour [15 miles per hour]). Observers will spend a minimum of 5 minutes at each survey location listening for calling frogs, but will terminate the survey early if frogs are detected.

To increase accuracy in the calculation of detectability of calling frogs, a small number of acoustic monitoring devices will be deployed at a subset of water bodies known to be occupied by frogs. Data from automated acoustic monitoring devices, which record calls throughout the day, will allow calculations of the probability of frogs calling on a given date, or at a specific time period and/or temperature range during the day, and will provide a direct estimate of the detectability of calling frogs.

Habitat and environmental characteristics (e.g., size and depth of water body, substrate, presence and type of emergent aquatic vegetation, distance to human disturbance, water quality [pH level, dissolved oxygen], ice cover, elevation, aspect, surrounding terrestrial vegetation, water and air temperature, precipitation, cloud cover, wind speed, time of day, beaver activity) will be recorded during the field surveys to facilitate the development of a Project-specific occupancy estimation model based on the habitat characteristics of the occupied water bodies. In addition, data from the vegetation and habitat mapping, wetland mapping, and wetland functional assessment studies (see Sections 11.5 and 11.7), and the literature (e.g., Stevens et al. 2006; AKNHP 2008) will be evaluated as potential model variables to characterize wood frog habitat.

With estimates of the detectability of wood frogs calculated from the field data collected for this study, the observed (“naïve”) occupancy rate of frogs in water bodies and wetlands will be corrected (to account for those frogs present but not detected) to produce a corrected occupancy rate for the water bodies and wetlands in each of the sampling regions.

#### *10.18.4.2. Bioassays for *Batrachochytrium dendrobatidis* (Bd)*

The specific assay and sampling methods for Bd will be determined through consultation with commercial or research laboratories. Currently available information indicates that no standard methods for bioassay of Bd have been proffered or certified by the U.S. Environmental Protection Agency (EPA) or other regulatory or standards agencies. The currently proposed strategy is to assess the presence/absence of Bd from swabs of frog skin, which would then be analyzed using a Polymerase Chain Reaction (PCR) technique to test for chytrid fungus.

Further consultation with the ADF&G and the U.S. Fish and Wildlife Service (USFWS) in early 2013 will be conducted to finalize the sampling protocol described here, but provisionally, frogs will be collected opportunistically during the field surveys with long-handled nets. The skin of the abdomen and/or foot webbing of each captured frog will be swabbed 25 times with a sterile



cotton swab, after which the frog will be released unharmed. The samples will then be sealed and refrigerated and analyzed later in the laboratory for the presence of chytrid DNA.

#### 10.18.4.3. Reporting and Data Deliverables

The reports and data deliverables for this study include the following:

- **Electronic copies of field data.** A geospatially-referenced relational database of field data collected during the 2013 and 2014 field seasons, including representative photographs of water body habitats occupied by wood frogs, will be prepared. Naming conventions of files and data fields, spatial resolution, map projections, and metadata descriptions will meet the data standards to be established for the Project.
- **Initial Study Report and Updated Study Report.** The Wood Frog Study results will be presented in the Initial and Updated Study reports, according to the schedule indicated in Table 10.18-1. The reports will include descriptions of the field methods, a map of the water bodies and wetlands surveyed, results of the occupancy surveys, and descriptions of the potential impacts to wood frogs from development of the Project.

#### 10.18.5. Consistency with Generally Accepted Scientific Practice

The Wood Frog Study will involve occupancy surveys of randomly selected water bodies and suitable wetland habitats, and will be conducted following currently accepted practices for the monitoring of amphibians, with field surveys designed to estimate detectability (USGS 2012). A similar occupancy survey of wood frogs in randomly selected water bodies was successfully conducted by ABR in 2007 on another large-scale project in southwest Alaska (see PLP 2011).

#### 10.18.6. Schedule

The Wood Frog Study is planned to be conducted over two years. The activities for each year are described in Table 10.18-1. Finalization of the sampling protocol for Bd will occur in February and March of 2013. Review of aerial imagery and selection of sampling regions and habitat areas to be surveyed will be conducted in March and April each year. Field surveys by a crew of two biologists will be conducted during the second and third weeks of May each year, with specific survey timing and duration to be determined annually, depending on snowmelt and lake-thaw information obtained from personnel working on other studies in the Project area each spring. Data analyses will be conducted during September–December each year. The Initial Study Report will be completed by February 2014 and the Updated Study Report will be completed by February 2015.

Technical Workgroup meetings will be planned on a quarterly basis in 2013 and 2014 to review study progress. Licensing participants will have the opportunity to review and comment on the Initial Study Report and Updated Study Report when they are completed.

#### 10.18.7. Relationship with Other Studies

As depicted in Figure 10.18-2, the Wood Frog Study will use information from the Vegetation and Wildlife Habitat Mapping Study in the Upper and Middle Susitna Basin (Section 11.5), the Wetland Mapping Study in the Upper and Middle Susitna Basin (Section 11.7), and the fish distribution and abundance studies in the Upper, Middle, and Lower Susitna River (Sections 9.5

and 9.6) to identify habitats potentially occupied by wood frogs. Potential habitats will be identified from air-photo interpretation during the mapping studies and, where available, from fish presence data in individual water bodies (those without fish being of higher value for frogs). Random sampling will be used to first select sampling regions for acoustic surveys of frogs from among all areas of suitable habitat in the study area, and then to select specific areas of habitat to be surveyed. Acoustical survey data from multiple visits will be used to estimate detectability of calling male frogs and to develop occupancy estimates for the areas of habitat surveyed. When completed, data from this study will be used in the Evaluation of Wildlife Habitat Use study (Section 10.19) to determine habitat values for wood frogs for each of the wildlife habitat types mapped for the Project.

The distribution information and habitat occupancy estimates for wood frogs determined in this study will be used to assess the potential impacts of the Project and to develop PM&E measures, as appropriate, during preparation of the FERC License Application in 2015.

Wood frogs potentially could be affected primarily by direct mortality during construction and by the loss of water bodies and wetlands suitable for breeding from the placement of fill and from inundation in the reservoir impoundment zone. Additional impacts could occur from the alteration of habitats due to erosion, fugitive dust accumulation, permafrost degradation, landslides, and off-road vehicle use. Aquatic habitats created by the impoundment may not be suitable for wood frogs due to their preference for smaller water bodies.

The impact assessment for wood frogs will be conducted by ascertaining which water bodies and wetland types are suitable for breeding using habitat characteristics that can be identified from aerial imagery interpretation, wildlife habitat and wetlands mapping, and fish survey data, as described above in Section 10.18.4.1. This information will allow the calculation of the amount of suitable habitat available before development. Using the corrected occupancy estimates from this study, the amount of the available habitat will be reduced to that amount most likely to be actually occupied (e.g., if the occupancy rate is estimated at 50 percent, then, on average, only 50 percent of the available habitat will be occupied). However, because all suitable habitat in the study area cannot be sampled, there will not be spatially explicit information for all sites to indicate which sites are actually occupied and which are not. With this available habitat and occupancy information, the Project footprint will be overlain, in GIS, on the map polygons representing suitable water body and wetland types to estimate the acreages of water bodies and wetlands that would be lost directly to fill or inundation. This acreage figure will then be reduced to account for the calculated occupancy rate, as noted above. The estimation of acreages of frog breeding habitats that could be affected by habitat alteration will be conducted similarly by overlaying habitat alteration buffers (surrounding the proposed Project infrastructure) to identify which habitats are likely to be affected by ancillary impacts associated with Project construction and operations. The size and number of habitat alteration buffer(s) to be used will be determined based upon the final specifications for Project construction and operations activities, which will be provided in the Project description.

Sampling for Bd in frogs in 2013 and 2014 will establish a baseline for comparison of the occurrence of Bd in frogs in the Project area after construction of the proposed Project.

Cumulative effects on wood frogs in the region of the proposed Project will be assessed in the FERC License Application document (to be prepared in 2015) and the details of that analysis (e.g., the spatial scale and temporal extent for cumulative effects) will be defined at that time.

### 10.18.8. Level of Effort and Cost

The Wood Frog Study is planned to be conducted over two years (2013–2014). A single field survey effort will be conducted each year in late spring (May) by a crew of two biologists. Based on previous occupancy surveys (PLP 2011), it is estimated that roughly 25 sites can be surveyed in a day. Occupancy surveys will be conducted for approximately 10 days each year. Helicopter support will be required for this study with multiple drop-offs and pick-ups in the afternoon and evening hours each day in the field (i.e., a dedicated helicopter may be required). The bulk of the costs associated with this study are for the field sampling, data analysis, and reporting. The projected cost for this study in each year is on the order of \$80,000, for an approximate estimated total of \$160,000 for both years.

### 10.18.9. Literature Cited

- ADF&G (Alaska Department of Fish and Game). 2006. *Our wealth maintained: A strategy for conserving Alaska's diverse wildlife and fish resources*. Alaska Department of Fish and Game Juneau. 824 pp.
- AEA (Alaska Energy Authority). 2012. Proposed Study Plan: Susitna-Watana Hydroelectric Project FERC Project No. 14241. July 2012. Prepared for the Federal Energy Regulatory Commission by the Alaska Energy Authority, Anchorage, Alaska.
- AKNHP (Alaska Natural Heritage Program). 2008. Alaska Wood Frog Monitoring Project results, 2002–2008. Available online: <http://aknhp.uaa.alaska.edu/zoology/citizen-science/alaska-wood-frog-monitoring/results-2002-2008/> (accessed October 2012).
- Anderson, B. C. 2004. An opportunistic amphibian inventory in Alaska's national parks, 2001–2003. Final report, National Park Service, Alaska Region Survey and Inventory Program, Anchorage. 44 pp.
- Babbitt, K. J., and G. W. Tanner. 1998. Effects of cover and predator size on survival and development of *Rana utricularia* tadpoles. *Oecologia* 114: 258–262.
- Blaustein, A. R., and D. B. Wake. 1990. Declining amphibian populations: a global phenomenon? *Trends in Ecology and Evolution* 5: 203–204.
- Cook, J. A., and S. O. MacDonald. 2003. Mammal inventory of Alaska's national parks and preserves: Denali National Park and Preserve. 2002 annual report for National Park Service, Alaska Region Survey and Inventory Program, Anchorage, by Idaho State University, Pocatello. 24 pp.
- France, R. L. 1997. The importance of beaver lodges in structuring littoral communities in boreal headwater lakes. *Canadian Journal of Zoology* 75: 1009–1013.
- Gotthardt, T. 2004. Monitoring the distribution of amphibians in the Cook Inlet watershed: 2003 final report. Alaska Natural Heritage Program, University of Alaska, Anchorage.
- Gotthardt, T. 2005. Wood frog conservation status report. Alaska Natural Heritage Program, University of Alaska, Anchorage.
- Hokit, D. G., and A. Brown. 2006. Distribution patterns of wood frogs (*Rana sylvatica*) in Denali National Park. *Northwestern Naturalist* 87: 128–137.

- MacDonald, S. O. 2010. *The amphibians and reptiles of Alaska: a field handbook*. Version 2.0. University of Alaska Museum, Fairbanks, and Museum of Southwestern Biology, Albuquerque, NM. Available online: <http://aknhp.uaa.alaska.edu/wp-content/uploads/2011/02/Herps-of-Alaska-Handbook-Final-Version-2-reduced.pdf> (accessed March 2012).
- MacKenzie, D. I., and J. A. Royle. 2005. Designing occupancy studies: general advice and allocating survey effort. *Journal of Applied Ecology* 42:1105–1114.
- McCallum, M. L. 2007. Amphibian decline or extinction? Current declines dwarf background extinction rate. *Journal of Herpetology* 41: 483–491.
- PLP (Pebble Limited Partnership). 2011. Pebble Project Environmental Baseline Document, 2004 through 2008. Chapter 16.12: Wood frogs—Mine Study Area. Pebble Limited Partnership, Anchorage. Available online: <http://www.pebbleresearch.com/> (accessed June 2012).
- Reeves, M. K. 2008. *Batrachochytrium dendrobatidis* in wood frogs (*Rana sylvatica*) from three national wildlife refuges in Alaska, USA. *Herpetological Review* 39: 68–70.
- Stevens, C. E., C. A. Paszkowski, and G. J. Scrimgeour. 2006. Older is better: Beaver ponds on boreal streams as breeding habitat for the wood frog. *Journal of Wildlife Management* 70: 1360–1371.
- USGS (U.S. Geological Survey). 2012. Amphibian Research and Monitoring Initiative. Available online: [http://armi.usgs.gov/amphibian\\_monitoring.php](http://armi.usgs.gov/amphibian_monitoring.php) (accessed October 2012).

**10.18.10. Tables**

**Table 10.18-1. Schedule for implementation of the Wood Frog Study.**

Activity	2013				2014				2015
	1 Q	2 Q	3 Q	4 Q	1 Q	2 Q	3 Q	4 Q	1Q
Finalize sampling protocol for Bd	—								
Review aerial imagery to select aquatic habitats to survey	—				—				
Field survey; survey timing and duration may be modified, depending on snowmelt and lake-thaw information obtained from other Project studies		—				—			
Data analysis			—	—			—	—	
Initial Study Report					△				
Updated Study Report									▲

Legend:

- Planned Activity
- △ Initial Study Report
- ▲ Updated Study Report

10.18.11. Figures




Figure 10.18-1. Wood frog study area.

### STUDY INTERDEPENDENCIES FOR WOOD FROG STUDY




Figure 10.18-2. Interdependencies for Wood Frog Study.

## **10.19. Evaluation of Wildlife Habitat Use**

### **10.19.1. General Description of the Proposed Study**

The Evaluation of Wildlife Habitat Use Study will be an analysis of both existing information on wildlife habitat use in Alaska (e.g., from the scientific literature) and new, Project-specific information on wildlife habitat use derived from survey data to be collected for the Project (see Sections 10.5 to 10.18). This habitat-use information will be used to systematically evaluate the use of the specific wildlife habitat types being mapped for the Project (see Section 11.5). In this study, categorical habitat values will be determined for each mapped habitat type and each wildlife species of concern to be assessed for impacts during the FERC licensing process.

#### **Study Goal and Objectives**

The goal of the study is to provide Project-specific habitat evaluation information for birds, mammals, and amphibians to facilitate quantitative assessments of the impacts on wildlife habitats from development of the proposed Project.

The wildlife habitat evaluation has two fundamental objectives:

- Use Project-specific survey data and the scientific literature to determine local habitat associations for those wildlife species occurring in the Project area that are of conservation, management, cultural, or ecological concern and that are specific to the wildlife habitat types to be mapped in the Project area.
- Categorically rank habitat values for each wildlife species of concern for each of the wildlife habitat types that will be mapped in the Project area.

The habitat-association data to be developed in this study, together with the wildlife habitats that will be mapped digitally in the Vegetation and Wildlife Habitat Mapping Study in the Upper and Middle Susitna Basin and the Riparian Vegetation Study Downstream of the Proposed Susitna-Watana Dam (see Sections 11.5 and 11.6, respectively), will be used in spatially-explicit analyses with a Geographic Information System (GIS) to derive quantitative estimates of habitat loss, habitat alteration, and disturbance effects for birds, mammals, and amphibians (see Section 10.19.7 below). This impact assessment work, which is not part of this study but is dependent on the results of this study, will be conducted during preparation of the FERC License Application for the Project.

### **10.19.2. Existing Information and Need for Additional Information**

Wildlife habitat evaluations for the Susitna basin were conducted in several studies in the early 1980s for the Alaska Power Authority (APA) Susitna Hydroelectric Project and for another study effort in the lower portions of the drainage (AEA 2011). Those habitat evaluations were based on vegetative cover types that were mapped within 16 kilometers (10 miles) on each side of the Susitna River between Gold Creek and the Maclaren River (TES 1982). That vegetation mapping and the subsequent habitat evaluations were conducted three decades ago.

Both the vegetation mapping and the habitat evaluations should be updated for the current Project for three primary reasons. First, the wildlife habitat evaluations completed in the early 1980s were based solely on vegetation types, not wildlife habitat types. Wildlife habitat maps provide land-cover classifications that are better suited to evaluations of habitat use by birds,



mammals, and amphibians than is a vegetation map alone, primarily through the incorporation of physiography, landform, and vegetation structure information (see Section 11.5). Second, populations of wildlife species undoubtedly have fluctuated in size since the early 1980s, and it is known that habitat use by birds and mammals can be influenced by density (a greater diversity of habitats often is used when densities are high). Third, vegetation cover, structure, and even landforms are likely to have changed to some degree within the Project area because of landslides, erosion, thermokarst, fire, forest succession, expansion/contraction/decadence of birch and aspen clones, and increases in woody shrub cover associated with increased summer temperatures. To provide accurate information to use in evaluating the impacts of habitat loss and alteration for wildlife species during the FERC licensing process, it is imperative that wildlife habitat evaluations be updated for the currently proposed Project, and that those habitat evaluations are based on a recently prepared wildlife habitat map for the Susitna basin.

### **10.19.3. Study Area**

The wildlife habitat evaluation study area will be identical to the area mapped for wildlife habitats in the upper and middle Susitna basin (Section 11.5), plus the area downstream of the proposed dam mapped to be mapped for riparian wildlife habitats (Section 11.6). These two areas overlap between the dam site and Gold Creek (Figure 10.19-1), but wildlife habitats in that section of the Susitna River floodplain will be mapped only in the Riparian Vegetation Study Downstream of the Proposed Susitna-Watana Dam (Section 11.6). The wildlife habitat evaluation study area (Figure 10.19-1) includes a 4-mile buffer surrounding those areas in the upper and middle Susitna basin that could be directly affected by Project construction and operations (the proposed reservoir impoundment zone, areas for infrastructure of the dam and powerhouse and supporting facilities, the proposed access route and transmission line corridors, and materials sites). The portion of the study area along the Susitna River downstream of Gold Creek includes the width of the active floodplain, as represented by the extent of riverine physiography (see Section 11.6). The downstream extent and width of the riparian zone to be evaluated in the Evaluation of Wildlife Habitat Use Study will match the final study area boundaries developed for the Riparian Vegetation Study Downstream of the Proposed Susitna-Watana Dam, which will be determined in the first quarter of 2013 (see Section 11.6).

### **10.19.4. Study Methods**

#### *10.19.4.1. Habitat Evaluation Procedures*

The proposed methods for the Evaluation of Wildlife Habitat Use Study involve the use of current and Project-specific survey data for birds, mammals, and amphibians in coordination and conjunction with the preparation of a current wildlife habitat map for the Project area. This study will be an office-based effort, performed after the wildlife habitat mapping for the Project area is completed. The methods to be used will follow those outlined in ABR (2008) and Schick and Davis (2008).

The first task in the Evaluation of Wildlife Habitat Use Study is selection of a set of wildlife species of concern for which Project-related habitat impacts will be evaluated. The selection criteria to be used to determine which animals are included will be finalized with input from the federal and state resource management agencies and other interested licensing participants in Q1 2013 as part of the planned Technical Workgroup (TWG) meetings, which will be scheduled

quarterly in 2013 (see Section 10.19.6 below). Specific criteria will be established for the species-selection process. It is proposed that a species be selected if it meets one or more of the following criteria:

- A federally- or state-protected species.
- A bird species of conservation and management concern, determined from lists maintained by various management agencies, agency working groups, and non-governmental conservation organizations (as outlined in the FERC–USFWS Memorandum of Understanding [MOU] on migratory birds; FERC and USFWS [2011]).
- A bird or mammal species of management concern for federal and/or state management agencies (primarily game and furbearer species).
- A species that is an important subsistence resource or is culturally significant for Alaska Natives.
- An ecologically important species with demonstrable ecosystem effects, such as ecosystem engineers (e.g., beaver), and species that occupy prominent positions in the trophic structure as predators or prey.

As agreed to during meetings with resource management agencies (see Appendices 3 and 4), the preliminary list of bird species of concern for the Project area (Table 10.19-1) comprises those species listed in Table 2 of the wildlife data-gap report for the Project (ABR 2011) and in Table 4.8-2 of the Project Pre-Application Document (PAD; AEA 2011), plus two additional shorebird species (Short-billed Dowitcher and Hudsonian Godwit) requested by the U.S. Fish and Wildlife Service (USFWS). The list of mammal species of concern will include big game, furbearers, and selected species of smaller mammals, including the little brown bat and Alaska tiny shrew. The list of wildlife species of concern, which is likely to include birds, mammals, and amphibians, will be refined further with input from resource management agencies.

A matrix will be constructed listing each species of concern and each wildlife habitat type mapped in the Project area, and a habitat-value ranking will be assigned to each cell in the matrix. As with the species selection process, the ranking procedure will be developed further with input from federal and state resource management agencies and other interested licensing participants, but it is likely that a habitat-value categorization system will be used (e.g., negligible, low, moderate, and high value). The habitat-value rankings will be derived in different ways among species, depending on the level of Project-specific data that are available to assess habitat use in each of the mapped wildlife habitat types. Observations of wildlife species will be tagged to mapped habitats using global positioning system (GPS) coordinates and a GIS, and the data quality will be assessed for each species and mapped habitat type (e.g., adequately sampled, under-sampled, or not sampled). Data-supported quantitative evaluations of habitat use will be employed whenever possible in the habitat-value rankings. However, in cases in which the habitats in question were under-sampled or not sampled, or for which sufficient Project-specific data are not available, then habitat-use information from the scientific literature and from field experience with the species elsewhere in Alaska will be used to derive habitat-value rankings.

Habitats will be ranked for the various life history stages of each of the species of concern addressed (e.g., breeding/calving, post-calving, spring and fall migration, overwintering) to encompass the complete seasonal range of habitat use. Additionally, specific habitat-use maps can be prepared for high-value game animals such as caribou, moose, and bears to illustrate

specific areas and seasons of use, in addition to identifying habitat types that are important to those species.

#### **10.19.4.2. Reporting and Deliverables**

Because the Evaluation of Wildlife Habitat Use Study cannot be completed until after the wildlife habitat mapping for the Project area is completed in October 2014, a brief Initial Study Report will be completed in February 2014, but the final habitat evaluations will not be available until the Updated Study Report is completed in February 2015 (see Section 10.19.6 below). The Updated Study Report will include descriptions of the methods used, including summaries of habitat use for each species assessed, and tables indicating habitat-values by species and habitat type. As agreed to with the resource management agencies (see Appendices 3 and 4), individual sections for each species assessed will be prepared in which the available habitat-use information will be linked to the specific habitat values derived (to illustrate the logic used in determining habitat values for each species).

#### **10.19.5. Consistency with Generally Accepted Scientific Practice**

The study methods discussed above have been successfully used for recent wildlife habitat evaluations on several projects in Alaska (e.g., ABR 2008; Schick and Davis 2008; PLP 2011). The methods have been favorably received by agency reviewers.

#### **10.19.6. Schedule**

The schedule for implementation of the Evaluation of Wildlife Habitat Use Study is summarized below (Table 10.19-2). The wildlife habitat evaluation can be completed only after the wildlife habitat mapping for the Project area is available in October 2014. Preliminary information to be used in the habitat-use rankings can be obtained through literature review in 2013 and earlier in 2014, however. The initial selection of species for analysis and accompanying literature review to support the habitat evaluations will be conducted during February–April 2013. A preliminary report of progress to date will be prepared for the Initial Study Report in February 2014 and the initial habitat-value rankings will be prepared during February–April 2014, using the preliminary results of wildlife field studies that are available by that time. The final selection of species for the final evaluation matrix will be completed by September 2014 and the final data analyses and habitat-value rankings will be conducted during September–December 2014, for presentation in the Updated Study Report in February 2015.

TWG meetings will be planned on a quarterly basis in 2013 and 2014 to review study progress. Licensing participants will have the opportunity to review and comment on the Initial Study Report and Updated Study Report.

#### **10.19.7. Relationship with Other Studies**

The relationships between the wildlife habitat-use evaluation and other Project studies are summarized here and illustrated below (Figure 10.19-2). Primary sources of information for the wildlife habitat-use evaluation include the wildlife habitat map polygons for the upper and middle Susitna basin from the Vegetation and Wildlife Habitat Mapping Study in the Upper and Middle Susitna Basin (Section 11.5), and the wildlife habitat map polygons for riparian areas downstream of the proposed dam from the Riparian Vegetation Mapping Study (Section 11.6).

As was described above, these mapped wildlife habitats will be evaluated for wildlife use and will be ranked categorically in terms of habitat value for a selected set of wildlife species of concern. Project-specific habitat-use information for mammals, birds, and amphibians will be obtained from each of the wildlife studies (Sections 10.5–10.18). These Project-specific data will be provided in GIS so that they can be directly associated with the mapped habitat types. From each of the wildlife studies, information on the locations of observations, the species and numbers recorded, seasonality, and behaviors observed, when available, will be used to evaluate the use of the wildlife habitats mapped for the Project.

The information on wildlife habitat values derived in this study will be used in the FERC License Application to assess the expected impacts of the proposed Project on the habitats known to be used by each wildlife species of concern in the study area. In addition, the wildlife habitat values will be used in the License Application to develop protection, mitigation, and enhancement (PM&E) measures, as appropriate.

Data from the Evaluation of Wildlife Habitat Use Study will be used in quantitative assessments of habitat loss and habitat alteration for each of the wildlife species of concern. With habitat-value rankings for each bird, mammal, and amphibian species of concern for each mapped habitat type, the areas within the Project footprint that are important for each species of concern can be identified, and the total areas of each to be directly affected (e.g., habitat loss and habitat alteration) by development of the Project can be determined quantitatively in GIS. Similarly, the indirect effects of disturbance will be assessed by applying species-specific disturbance buffers to the Project footprint and determining quantitatively the total areas of important habitats for each species of concern that could be influenced indirectly by disturbance effects during Project construction and operations. Data from the Evaluation of Wildlife Habitat Use Study will also be used to help address the potential for fragmentation of habitat patches for species of concern because of Project development.

Also in the FERC License Application, for areas downstream of the proposed dam, the habitat-value rankings from this study will be used to help predict how wildlife species will respond to the changes in riparian wildlife habitats in the Susitna River floodplain that are expected to occur with construction and operation of the proposed dam.

As agreed to during meetings with resource management agencies (see Appendices 3 and 4), the finer-scale habitat types mapped in the Project area (see Section 11.5) will be “crosswalked” with the coarser-scale habitats (30-meter pixel resolution) mapped in the Alaska Gap Analysis Project (GAP). The habitat-value rankings for each wildlife species of concern in each mapped habitat type in the Project area will also be “crosswalked” to the coarser-scale GAP habitats, and averaged, when multiple values need to be combined, to derive appropriately-scaled habitat rankings. With the habitat-value rankings upgraded to the GAP habitat types, the habitat loss and habitat alteration effects from the proposed Project can be placed in a broader regional context (e.g., habitat impacts can be assessed at the eco-regional scale).

#### **10.19.8. Level of Effort and Cost**

The wildlife habitat evaluation will be an office-based effort and is expected to be completed relatively quickly once the wildlife habitat mapping tasks are completed. The Evaluation of Wildlife Habitat Use Study can be completed in several months. The habitat evaluation will be conducted by up to two vegetation ecologists and four wildlife biologists (with specific expertise

with various vertebrate species groups). The total cost of this study over both years is estimated to be approximately \$200,000.

### 10.19.9. Literature Cited

- ABR. 2008. Chuitna Coal Project: Wildlife Protection Plan, Part D7-2. Final report prepared for Mine Engineers, Inc., Cheyenne, WY, on behalf of PacRim Coal LP, Anchorage, by ABR, Inc. —Environmental Research & Services, Anchorage. 153 pp.
- ABR. 2011. Wildlife data-gap analysis for the proposed Susitna–Watana Hydroelectric Project. Draft report, August 16, 2011. Report for the Alaska Energy Authority by ABR, Inc.—Environmental Research and Services, Fairbanks. 114 pp.
- ADF&G. 2006. *Our wealth maintained: A strategy for conserving Alaska’s diverse wildlife and fish resources*. Alaska Department of Fish and Game, Juneau.
- AEA. 2011. Pre-application Document, Susitna–Watana Hydroelectric Project, FERC Project No. 14241. Volume I, Section 4.6: Wildlife and Botanical Resources.
- APA (Alaska Power Authority). 1985. Before the Federal Energy Regulatory Commission: Draft amended application for license for major project—Susitna Hydroelectric Project. Volume 11, Exhibit E, Chapter 3: Fish, wildlife, and botanical resources.
- ASG (Alaska Shorebird Group). 2008. Alaska Shorebird Conservation Plan. Version II. Anchorage.
- BLM (Bureau of Land Management). 2010a. BLM–Alaska sensitive animal and plant lists. Alaska State Office, Anchorage.
- BLM. 2010b. BLM–Alaska watch lists. Alaska State Office, Anchorage.
- BPIFWG (Boreal Partners in Flight Working Group). 1999. Landbird conservation plan for Alaska biogeographic regions. Version 1.0. U.S. Fish and Wildlife Service, Anchorage.
- FERC and USFWS. 2011. Memorandum of Understanding between the Federal Energy Regulatory Commission and the U.S. Department of the Interior United States Fish and Wildlife Service Regarding Implementation of Executive Order 13186, “Responsibilities of Federal Agencies to protect Migratory Birds.” March 2011. 13 pp.
- Kessel, B., S. O. MacDonald, D. D. Gibson, B. A. Cooper, and B. A. Anderson. 1982. Susitna Hydroelectric Project environmental studies, Phase I final report—Subtask 7.11: Birds and non-game mammals. Report prepared by University of Alaska Museum, Fairbanks, and Terrestrial Environmental Specialists, Inc., Phoenix, NY, for Alaska Power Authority, Anchorage. 149 pp.
- Kushlan, J. A., M. J. Steinkamp, K. C. Parsons, J. Capp., M. Acosta Cruz, M. Coulter, I. Davidson, L. Dickson, N. Edelson, R. Elliot, R. M. Erwin, S. Hatch, S. Kress, R. Milko, S. Miller, K. Mills, R. Paul, R. Phillips, J. E. Saliva, B. Sydeman, J. Trapp, J. Wheeler, and K. Wohl. 2002. *Waterbird conservation for the Americas: the North American Waterbird Conservation Plan*. Version 1. Waterbird Conservation for the Americas, Washington, D.C. 78 pp.
- Kushlan, J. A., M. J. Steinkamp, K. C. Parsons, J. Capp., M. Acosta Cruz, M. Coulter, I. Davidson, L. Dickson, N. Edelson, R. Elliot, R. M. Erwin, S. Hatch, S. Kress, R. Milko,

- S. Miller, K. Mills, R. Paul, R. Phillips, J. E. Saliva, B. Sydeman, J. Trapp, J. Wheeler, and K. Wohl. 2006. Conservation status assessment factor scores and categories of concern for solitary-nesting waterbird species. Addendum to *Waterbird conservation for the Americas: North American Waterbird Conservation Plan*. Version 1, April 17, 2006. Waterbird Conservation for the Americas, Washington, D.C.
- North American Waterfowl Management Plan Committee. 2004. North American Waterfowl Management Plan, 2004 — Implementation framework: strengthening the biological foundation. Canadian Wildlife Service, U.S. Fish and Wildlife Service, Secretaria de Medio Ambiente y Recursos Naturales. 106 pp.
- PLP (Pebble Limited Partnership). 2011. Pebble Project Environmental Baseline Document, 2004 through 2008. Pebble Limited Partnership, Anchorage. Available online: <http://www.pebbleresearch.com/> (accessed June 16, 2012).
- Schick, C. T., and W. A. Davis. 2008. Wildlife habitat mapping and evaluation of habitat use by wildlife at the Stewart River Training Area, Alaska. Final report, prepared for Alaska Army National Guard, Fort Richardson, by ABR, Inc.—Environmental Research & Services, Anchorage. 54 pp.
- TES (Terrestrial Environmental Specialists). 1982. Susitna Hydroelectric Project, Task 7— Environmental studies, wildlife ecology: wildlife habitat-value analysis. Report prepared by Terrestrial Environmental Specialists, Inc., Phoenix, NY, for Acres American, Inc., Buffalo, NY. 100 pp.
- USFWS (U.S. Fish and Wildlife Service). 2008. Birds of Conservation Concern, 2008. Division of Migratory Bird Management, Arlington, VA. 85 pp. Available online (accessed July 12, 2011): <http://www.fws.gov/migratorybirds>.
- USFWS. 2009. Birds of Management Concern. Division of Migratory Bird Management, Arlington, VA. Available online (accessed July 12, 2011): [http://www.fws.gov/migratorybirds/NewReportsPublications/SpecialTopics/BirdsofManagementConcern09\[1\].pdf](http://www.fws.gov/migratorybirds/NewReportsPublications/SpecialTopics/BirdsofManagementConcern09[1].pdf).

**10.19.10. Tables**

**Table 10.19-1. Bird species of conservation/management concern that are known or likely to occur in the Susitna River basin, Alaska.**

English Name	USFWS BCC <sup>1</sup>	USFWS BMC <sup>2</sup>	ADFG <sup>3</sup>	BLM <sup>4</sup>	NAWCP <sup>5</sup>	NAWMP <sup>6</sup>	ASG (USSCP) <sup>7</sup>	BPIF (PIF) <sup>8</sup>
Greater White-fronted Goose (Tule)		■				■		
Snow Goose		■						
Brant		■				■		
Canada Goose		■				■		
Trumpeter Swan		■		■				
Tundra Swan		■						
Gadwall		■						
American Wigeon		■				■		
Mallard		■				■		
Blue-winged Teal		■				■		
Northern Shoveler		■						
Northern Pintail		■				■		
Green-winged Teal		■						
Canvasback		■				■		
Redhead		■				■		
Ring-necked Duck		■						
Greater Scaup		■						
Lesser Scaup		■				■		
Harlequin Duck		■						
Surf Scoter		■	■			■		
White-winged Scoter		■	■			■		
Black Scoter		■				■		
Long-tailed Duck		■				■		
Common Goldeneye		■				■		
Rock Ptarmigan			■					

**Table 10.19-1. Continued.**

English Name	USFWS BCC <sup>1</sup>	USFWS BMC <sup>2</sup>	ADFG <sup>3</sup>	BLM <sup>4</sup>	NAWCP <sup>5</sup>	NAWMP <sup>6</sup>	ASG (USSCP) <sup>7</sup>	BPIF (PIF) <sup>8</sup>
White-tailed Ptarmigan								■
Red-throated Loon	■		■	■*	■			
Pacific Loon			■					
Common Loon			■					
Horned Grebe	■		■		■			
Red-necked Grebe			■					
Osprey			■					
Bald Eagle			■					
Northern Harrier			■					
Sharp-shinned Hawk			■					
Northern Goshawk			■					
Red-tailed Hawk			■					
Golden Eagle			■	■				
Merlin			■					
Gyrfalcon			■					■
Peregrine Falcon <sup>9</sup>	■		■					
American Golden-Plover							■	
Solitary Sandpiper	■	■	■				■	
Lesser Yellowlegs	■	■	■				■	
Upland Sandpiper	■	■					■	
Whimbrel	■	■					■	
Hudsonian Godwit	■	■					■	
Ruddy Turnstone <sup>10</sup>							■	
Black Turnstone <sup>10</sup>							■	
Short-billed Dowitcher	■	■					■	
Surfbird							■	
Sanderling							■	



**Table 10.19-1. Continued.**

English Name	USFWS BCC <sup>1</sup>	USFWS BMC <sup>2</sup>	ADFG <sup>3</sup>	BLM <sup>4</sup>	NAWCP <sup>5</sup>	NAWMP <sup>6</sup>	ASG (USSCP) <sup>7</sup>	BPIF (PIF) <sup>8</sup>
Wilson's Snipe		■						
Black-legged Kittiwake			■					
Arctic Tern			■					
Great Horned Owl			■					
Snowy Owl			■					
Northern Hawk Owl			■					
Short-eared Owl	■		■	■				■
Boreal Owl			■					■
Belted Kingfisher			■					
Hairy Woodpecker			■					
American Three-toed Woodpecker			■					
Black-backed Woodpecker			■					■
Northern Flicker			■					
Olive-sided Flycatcher	■		■	■				■
Western Wood-Pewee								■
Northern Shrike								■
Violet-green Swallow			■					
Bank Swallow			■					
Cliff Swallow			■					
Boreal Chickadee			■					
Brown Creeper			■					
American Dipper								■
Golden-crowned Kinglet			■					
Gray-cheeked Thrush				■ *				■
Hermit Thrush			■					
Varied Thrush			■					■
Bohemian Waxwing								■

**Table 10.19-1. Continued.**

English Name	USFWS BCC <sup>1</sup>	USFWS BMC <sup>2</sup>	ADFG <sup>3</sup>	BLM <sup>4</sup>	NAWCP <sup>5</sup>	NAWMP <sup>6</sup>	ASG (USSCP) <sup>7</sup>	BPIF (PIF) <sup>8</sup>
Smith's Longspur	■		■					■
Blackpoll Warbler			■	■				■
Townsend's Warbler			■	■*				■
Wilson's Warbler			■					
White-crowned Sparrow			■					
Golden-crowned Sparrow								■
Dark-eyed Junco			■					
Rusty Blackbird	■		■	■				■
Gray-crowned Rosy Finch			■					
Pine Grosbeak			■					
White-winged Crossbill			■					■
Pine Siskin			■					

Species list derived from Kessel et al. (1982) and APA (1985: Appendices E5.3 and E6.3), plus Townsend's Warbler, Hudsonian Godwit, and Short-billed Dowitcher.

<sup>1</sup> USFWS (2008) Birds of Conservation Concern.

<sup>2</sup> USFWS (2009) Birds of Management Concern.

<sup>3</sup> ADF&G (2006) Featured Species.

<sup>4</sup> BLM (2010a) Sensitive Species; asterisk denotes Watch List Species (BLM 2010b).

<sup>5</sup> North American Waterbird Conservation Plan (Kushlan et al. 2002, 2006).

<sup>6</sup> North American Waterfowl Management Plan Committee (2004).

<sup>7</sup> Alaska Shorebird Group (2008).

<sup>8</sup> Boreal Partners in Flight Working Group (1999).

<sup>9</sup> Previously listed as threatened under the Endangered Species Act (ESA), the American Peregrine Falcon (*Falco peregrinus anatum*) was delisted in August 1999.

<sup>10</sup> Species identity (Ruddy Turnstone, Black Turnstone) of sole record in the Susitna basin was unconfirmed (Kessel et al. 1982), but both are on the ASG list.

**Table 10.19-2. Schedule for implementation of the wildlife habitat-use evaluation.**

Activity	2013				2014				2015
	1 Q	2 Q	3 Q	4 Q	1 Q	2 Q	3 Q	4 Q	1 Q
Initial selection of species for analysis	—								
Literature review of habitat-use information	—	—							
Initial Study Report					△				
Initial habitat-value ranking					—				
Final selection of species for analysis								—	
Data analysis and habitat-value ranking								—	—
Updated Study Report									▲

Legend:

- Planned Activity
- △ Initial Study Report
- ▲ Updated Study Report

10.19.11. Figures




Figure 10.19-1. Study area for evaluation of wildlife habitat use. The study area is a combination of the wildlife habitat mapping areas from the Vegetation and Habitat Mapping Study (Section 11.5) and the Riparian Vegetation Study (Section 11.6).

### STUDY INTERDEPENDENCIES FOR WILDLIFE HABITAT-USE EVALUATION




Figure 10.19-2. Study interdependencies for the wildlife habitat-use evaluation.

## 10.20. Wildlife Harvest Analysis

### 10.20.1. General Description of the Proposed Study

The wildlife harvest analysis study is an office-based study of Alaska Department of Fish and Game (ADF&G) and U.S. Fish and Wildlife (USFWS) harvest records for large mammals and furbearers, as well as for smaller mammals and upland gamebirds (if data are available). In this study, AEA will characterize past and current hunter effort and harvest levels in the region of the proposed Project by summarizing and analyzing data from the ADF&G harvest database for Alaska, which also includes some harvest data from subsistence users reported to USFWS.

#### Study Goal and Objectives

Construction and operation of the Project will alter human access to the region through construction of the access road and power transmission corridors, and through the creation of the reservoir. Much of Alaska Game Management Unit (GMU) 13, which encompasses the Project area, is readily accessible by road and provides hunting opportunities for many Alaskans. Creating access points to the Project site from the Denali Highway to the north or from the rail corridor to the west may result in increased motorized vehicle access for hunters and recreational users to portions of GMU 13 that are currently remote. The potential for increased human access and activity within GMU Subunits 13A and 13E without additional understanding of the implications for game populations has been identified as a resource management concern by ADF&G.

The goal of this study is to compile and analyze information on the distribution of big game, furbearers, and small game (including both small mammals and upland gamebirds, assuming data are available) and to understand patterns of hunting effort and harvest in the study area. These data will provide information on identification of past and current trends in hunter access modes, hunting locations, and harvest locations, and identify potential Project-induced changes that are likely to alter hunter access or harvest patterns. These findings will help predict the impacts of those changes on wildlife harvests. This study is a multi-year effort that began in 2012 (AEA 2012).

Specifically, this study has three primary objectives:

- Identify past and current harvest effort for large and small game including furbearers, harvest locations, access modes and routes.
- Compare current harvest locations of large and small game, including furbearers, with data on the seasonal distribution, abundance, and movements of harvested species, using the results of other, concurrent Project studies on big game and furbearers (Sections 10.5–10.11).
- Provide harvest data for use in the analyses to be conducted for the recreation and subsistence resource studies (Sections 12.5 and 14.5, respectively).

The information developed in this study will be used to help develop any necessary measures to address Project impacts on hunting opportunities, hunter distribution, and impacts to game species abundance.

### **10.20.2. Existing Information and Need for Additional Information**

The wildlife data-gap analysis conducted for the Project (ABR 2011) identified the need for an updated drainage-specific compilation of subsistence, sport hunter, and trapper harvest data for big game and furbearers. Hunter access to this region has changed since the 1980s, but potential changes in patterns of harvest at this scale have not been evaluated or compared with distribution of harvested species. Compilation of historic data may be useful for identifying trends in human access and harvest locations over the past decades and will provide information that may inform ADF&G's management goals for big game and furbearers in the Project area.

ADF&G documents legal sport hunting and trapping in Alaska through the collection of harvest reports and sealing records of hides for certain furbearers. Harvest reports are required to be submitted by hunters for some big game species. Hunting effort and harvest success are summarized from harvest reports and sealing records by GMU, subunit, and, when possible, by smaller Uniform Coding Units (UCU) that are delineated based on watersheds at a sub-basin level. These data are compiled and stored by ADF&G in a statewide harvest database. In addition, a trapper questionnaire is issued annually to compile trappers' views of various wildlife species in their areas (Schumacher 2010) and some subsistence hunting activity is summarized based on household surveys. Information on harvest as a part of federal subsistence hunts on federal land is maintained by USFWS and will need to be obtained through a separate data sharing agreement.

This information from ADF&G is available to be summarized and analyzed to elucidate spatial and temporal patterns of hunting effort and harvest success. It also provides some information on access types, use of guides, and residency of hunters. These data can be compared with data on the distribution of game mammals and the analyses can be used to help predict the impact of the Project on hunting opportunities and hunter distribution, and impacts on game mammals. Subsistence surveys will be conducted by ADF&G to gather current information for communities near the Project area. Additional information on subsistence harvests will also be available from other studies.

The following issues identified in the Pre-Application Document (PAD) (AEA 2011) will be addressed in this study:

- W4: Potential impact of changes in predator and prey abundance and distribution related to increased human activities and habitat changes resulting from Project development.
- W5: Potential impacts to wildlife from changes in hunting, vehicular use, noise, and other disturbances due to increased human presence resulting from Project development.

### **10.20.3. Study Area**

The study area (Figure 10.20-1) includes GMU Subunits 13A, 13B, 13E, 14B, 16A, and portions of 20A. These GMUs were selected because hunting and trapping activities in portions of each of these GMUs may be influenced directly or indirectly by Project construction and operations, including the reservoir inundation zone, associated facility sites, laydown/storage areas, and access road and power transmission corridors. The study area is based on GMUs conforming with the harvest data available (which is recorded by GMU) and because hunting and trapping in the region of the Project is managed by GMU.

#### **10.20.4. Study Methods**

In this study, AEA will use existing data, as well as new data to be collected during concurrent studies, to assess the spatial and temporal patterns and success of hunting and trapping efforts and to examine relationships between effort, harvest, and the distribution of wildlife, as indicated by telemetry studies and other surveys. Existing data from harvest reports will be compiled and reviewed to assess their adequacy to address Project-related changes in human access. These data will be shared with researchers conducting the recreation and subsistence resource studies (Sections 12.5 and 14.5). The methods used in this study will include the following tasks:

- Compilation and analysis of ADF&G harvest database records
- Review of ADF&G management reports
- Review of ADF&G trapper questionnaires
- Review of ADF&G small game outlook and harvest surveys
- Review of ADF&G and USFWS subsistence surveys and harvest reports
- Interviews with regional biologists
- Comparison of harvest patterns with development plans and the distribution of game mammals and birds

Initial efforts will focus on compilation and analysis of hunter effort and harvest success within harvest report units contained within the ADF&G harvest-record database. The spatial resolution, adequacy, and completeness of the harvest data record for detecting potential changes in use of wildlife resources in the Project area will be evaluated.

The study will build on results of the wildlife harvest data analysis begun in 2012 and will incorporate new harvest data as they become available, as well as the results of the ADF&G moose, caribou, and ptarmigan telemetry studies begun in 2012. Harvest patterns will be compared with seasonal distribution and movements revealed by the telemetry data on moose, caribou, and ptarmigan.

A relational database of harvest and effort data used in the analysis will be prepared. Naming conventions of files, data fields, and metadata descriptions will meet the Alaska Department of Natural Resources (ADNR) standards established for the Project. Harvest effort and success will be calculated at the highest spatial resolution possible given the quality of the data (GMUs, Subunits, or UCUs) and compared with the best available estimates of game populations, hunting regulations, and access. Hunter effort and harvest success maps showing big game and furbearer species will be developed based on the relational database developed from the ADF&G harvest database. All map and spatial data products will be delivered in the two-dimensional Alaska Albers Conical Equal Area projection, and North American Datum of 1983 (NAD 83) horizontal datum, consistent with ADNR standards.

#### **10.20.5. Consistency with Generally Accepted Scientific Practices**

Harvest data will be analyzed according to commonly accepted statistical techniques. Spatial statistics will be conducted with commonly accepted techniques such as fixed-kernel density estimation with least-squares cross validation or plug-in bandwidth selection (Seaman and Powell 1996; Gitzen et al. 2006).



### **10.20.6. Schedule**

This study is a multi-year effort that began in 2012 with data transfers from ADF&G and USFWS for dates from 2011 and earlier (AEA 2012). The schedule planned for 2013–2014 activities is depicted in Table 10.20-1. Transfer of 2012 harvest and subsistence data from ADF&G and USFWS is planned for July 2013 and 2014, depending on the availability of summarized data from their geodatabases (data transfer may occur somewhat later in the third quarter [3Q]). The data received from those agencies will be compiled into the Project-specific geodatabase for summary and analysis in the fourth quarter [4Q] each year, which will be used in the Initial Study Report to be completed by February 2014, and in the Updated Study Report to be completed by February 2015. Updates on the study progress will be provided during Technical Workgroup meetings which will be held quarterly in 2013 and 2014.

### **10.20.7. Relationship with Other Studies**

As depicted in Figure 10.20-2, data inputs for the wildlife harvest analysis will be required annually from the harvest databases maintained by state (ADF&G) and federal (USFWS) agencies. Those data will be compiled into a Project-specific geodatabase of harvest data, organized by species, date, method, and location (reporting area), which will be used to prepare spatially-explicit summaries of harvests in various portions of the study area to as fine a level of spatial resolution as is supported by the data. The data outputs from these analyses will be provided to the Subsistence Resource Study (Section 14.5) and the Recreation Resource Study (Section 12.5) so that subsistence and sport harvests, respectively, can be evaluated and compared with other human uses of the Project area.

During the impact assessment that will be conducted for the FERC License Application in 2015, the results of the wildlife harvest analysis will be used both directly and indirectly (through the other studies mentioned in the preceding paragraph) in the assessment of impacts and in the identification of any appropriate protection, mitigation, and enhancement (PM&E) measures. Data on the recent and current distribution of harvest effort and harvest success in the study area will be used to assess potential Project impacts on hunting and trapping effort and harvest success. The assessment of impacts on hunting and trapping effort and harvest success will be coordinated with the Recreational Resources Study and the Subsistence Resources Study (Sections 12.5 and 14.5, respectively) to assess how the expected changes in land use and access in the Project area may affect patterns of hunting and trapping. The direct and indirect impacts of the Project on game animal populations will be assessed in other wildlife studies (Sections 10.5–10.11 and 10.17) by conducting geospatial analyses using information on the responses of the species to other development projects, as documented in the scientific literature. Those geospatial analyses will overlay the Project footprint and species-specific habitat alteration and disturbance buffers on the known locations of use by the species of interest, as determined from Project-specific survey data and the Evaluation of Wildlife Habitat Use (Section 10.19). Similarly, Geographic Information System (GIS) analyses of potential impacts on hunting and trapping effort and harvest success will be conducted for this study by overlaying the Project footprint and species-specific habitat alteration and disturbance buffers on the known locations of harvest data obtained in this study.

### **10.20.8. Level of Effort and Cost**

This study will focus on analyzing existing harvest data and new data collected for other wildlife, subsistence, and recreational studies to maximize the information gained from these data. Thus, basic questions associated with human harvest of game animals in and near the Project area can be analyzed in a cost-effective manner. The estimated total cost of the study is less than \$100,000 over both years.

### **10.20.9. Literature Cited**

- ABR. 2011. Wildlife data-gap analysis for the proposed Susitna–Watana Hydroelectric Project. Draft report, August 16, 2011. Report for the Alaska Energy Authority by ABR, Inc.—Environmental Research and Services, Fairbanks, Alaska. 114 pp.
- AEA. 2011. Pre-Application Document: Susitna–Watana Hydroelectric Project FERC Project No. 14241. December 2011. Prepared for the Federal Energy Regulatory Commission, Washington, DC.
- AEA. 2012. Past and current big game and furbearer harvest study for the Susitna–Watana Hydroelectric Project, FERC Project No. 14241. Draft final version (March 21, 2012). Alaska Energy Authority, Anchorage.
- Gitzen, R. A., J. J. Millspaugh, and B. J. Kernohan. 2006. Bandwidth selection for fixed-kernel analysis of animal utilization distributions. *Journal of Wildlife Management* 70: 1334–1344.
- Schumacher, T. 2010. Trapper questionnaire: Statewide annual report, 1 July 2008–30 June 2009. Alaska Department of Fish and Game, Division of Wildlife Conservation, Juneau.
- Seaman, D. E., and R. A. Powell. 1996. An evaluation of the accuracy of kernel density estimators for home range analysis. *Ecology* 77: 2075–2085.

**10.20.10. Tables**

**Table 10.20-1. Schedule for implementation of the Wildlife Harvest Analysis.**

Activity	2012				2013				2014				2015
	1 Q	2 Q	3 Q	4 Q	1 Q	2 Q	3 Q	4 Q	1 Q	2 Q	3 Q	4 Q	1 Q
Transfer of 2012 harvest data							—						
Analysis of 2012 harvest data and preparation of Initial Study Report, to be completed in February 2014								—	△				
Transfer of 2013 harvest data											—		
Analysis of 2013 harvest data and preparation of Updated Study Report, to be completed in February 2015												—	▲

Legend:

- Planned Activity
- △ Initial Study Report
- ▲ Updated Study Report

10.20.11. Figures




Figure 10.20-1. Study area for the Wildlife Harvest Analysis.

## STUDY INTERDEPENDENCIES FOR WILDLIFE HARVEST ANALYSIS




Figure 10.20-2. Study interdependencies for the Wildlife Harvest Analysis.