



SUSITNA-WATANA HYDRO


Clean, reliable energy for the next 100 years.

ALASKA ENERGY AUTHORITY

AEA11-022

16-1401-TM-012014


Figures




- Explanation**
- Quaternary fault, solid where well constrained, long dashed line where moderately constrained, short dashed line where inferred (Alaska Division of Geological and Geophysical Surveys, 2012)
 - Field work planned in 2013 based on results of TM-8 (FCL, 2013)
 - No field work planned in 2013 based on results of TM-8 (FCL, 2013)
 - ★ Proposed Watana site

DRAFT

79_218900_Alaska_Railbelt/2189_Lineament Report October 2013, modified 12.02.13





DRAFT



- Explanation**
- Quaternary fault, solid where well constrained, long dash where moderately constrained, short dash where inferred (Koehler et al., 2012)
 - ★ Proposed Watana site
- GPS Tracks
(by reconnaissance date)**
- | | |
|-----------|-----------|
| 7/11/2013 | 7/19/2013 |
| 7/12/2013 | 7/21/2013 |
| 7/13/2013 | 7/22/2013 |
| 7/14/2013 | 7/23/2013 |
| 7/15/2013 | 7/24/2013 |
| 7/16/2013 | 9/4/2013 |
| 7/17/2013 | 9/5/2013 |
| 7/18/2013 | |


Coordinates on NAD83 UTM 6 North.
Elevation from INSAR data and USGS SRTM data.

79_218900_Alaska_Railbelt/2189_Lineament Report October 2013, modified 12.02.13




Notes: 1. See Figures A0.2, A0.3, A0.4, and A0.5 for explanation.
 2. Geology by Wilson et al., 1998


DRAFT



View looking northeast from location A towards the confluence of the Jack River and the East Fork Jack River. Arrows point along the alignment of mapped lineaments. Note absence of linear expression in Quaternary deposits.



View looking southwest from location B along alignment of linear features. Arrows indicate the alignment of the mapped lineaments.



View looking southwest from location C at a detailed view of aligned uphill-facing scarps. Note Thf contact is up-slope from the scarp in the distance.

79_218900_Alaska_Railbelt/2189_Lineament Report October 2013, modified 01.06.14

Attributes of lineaments mapped by FCL (2013) that apply to all figures and plates in Appendix A

Reconnaissance (INSAR)

- 1 - 5
- 10
- 77
- 88

Detail (LiDAR)

- 1 - 5
- 10
- 77
- 88

Lineament Groups

- 17a Lineament group mapped for this study coinciding with previously mapped fault or lineament
- 25 No previously mapped fault or lineament coincides with lineament group

Attribute	Cross Section Morphology*	Description	Examples
1		Linear break-in-slope bisecting a planar surface	Uphill- or downhill-facing scarps, ateral moraines or kame deposits along lateral margins of valley glaciers
2		Abrupt changes in slope adjacent to otherwise relatively horizontal (and planar) surfaces	Linear range fronts, faceted ridges, terrace risers, steep downstream faces of roushe mountonees
3		Linear U-shaped trough	Glacial valleys, ice-scoured flutes, flood-scoured flutes,
4		Linear V-shaped trough	Active stream channels
5		Linear ridges	Drumlins, water-scoured terrain, eskers
6 (also 77)	n/a	A series of aligned features	Could include attributes #1 -5 above and/or aligned saddles, tonal lineaments, etc.
66	n/a	Data artifacts	Linear seams between data sets collected on different dates
88	n/a	A series of aligned features, which are too small to individually map at the given scale	Could include features with attributes #1-5 above and/or aligned saddles, tonal lineaments, etc.
99	n/a	A line which encloses a broad expanse of features all having the same orientation	An area of jointing or of glacial striae all having the same, parallel orientation
10	n/a	Anthropogenic lineaments	Roads, rail roads, power lines and other linear clearings, etc.

Notes: *Arrow points to location of the mapped feature.

Explanation for relevant geologic units of Williams and Galloway (1986) shown on Figure A20.5 and A23.1

Geologic Units


Bottom deposits of 914 - 975 m lake
Overprint denoting glacial drift that is mantled by bottom sediments of glacial lake that extended to 914 - 975 m above modern sea level, largely confined to middle Susitna valley, above ice dam below Fog Lake (off map) and apparently bounded on east and south side by glacier ice. Does not cover late(st) Wisconsin (last major) morainial systems. No shoreline features are mapped.

Bottom deposits intermediate (777 - 747) lake
Overprint denoting bottom deposits of a local lake that covered melting glacier ice between Tyone Lake and Lake Louise, apparently behind Tyone Spillway, and drained as the elevation of the spillway was cut down from 777 m to 747 m above sea level while stagnant ice was still in valley bottom.

Bottom deposits of last regional lake
Overprint denoting drape of bottom deposits over drift and thick lake sediments that persisted in Copper River drainage basin from just before deposition of Old Man moraines to a time when glaciers had retreated to within 16 to 24 km of present glaciers: older than 13,000 years.


Symbols

- Location and letter designation of radiocarbon-dated stratigraphic section in accompanying text.
- Ice boundary, morainial ridge, kame terrace, delta, or other ice contact feature marking edge of glacier: hachures toward glacier.
- Shoreline of regional lake: mapped for the lake in Copper River basin where at 747 m (maximum elevation); the elevation to which Tyone Spillway was eroded, and successively lower levels in the northern part of area between 747 m and 701 m above sea level. Lesser recessional shorelines mapped by Nichols and Yehle (1969) not shown.
- Upper limit of post-glacial (Holocene, in part) shoreline of Tazlina Lake from elevation 564 m down to present lake level 544 m caused by lowering of lake as Tazlina River has deepened its canyon.
- Delta of glacial lake, including those of modern glacial lakes such as Tazlina Lake.
- Linear or drumlinoid feature, due to ice scour, direction of ice movement indicated by arrow.
- Spillway for glacial meltwater, including that stored in large glacial lakes.
- Contact between map units where not glacial boundary, most commonly between different levels of lake deposits.
- Active (?) fault, lower Sonoma Creek, offsetting unconsolidated deposits.
- Location of selected erratic boulders, mountain top erratic stones transported by glaciers, e.g. Sheep Mountain; many occurrences on mountains lower than 1829 m not shown.



Base data from ASTER Global Digital Elevation Model (ASTER GDEM is a product of METI and NASA)

Explanation




Extent of
INSAR data

Extent of LiDAR Data

- Area A
- Area B
- Area C
- Area D

Note: Extent of Landsat imagery and
ASTER GDEM elevation data
are greater than the area shown
in figure.



DRAFT



Date 01/06/14



SUSITNA-WATANA HYDROELECTRIC PROJECT

EXTENT OF GEOSPATIAL DATA


FIGURE

2-5



From Gray, 2001.

Note linearity of channels, lack of contributing watershed area, and steep sidewalls.



From <http://www.landforms.ca/cairngarms/meltwater%20channels.htm>, last accessed 1 October, 2013.

These sub-ice channels are cut through interfluves, seen as notches on the skyline.




Date 10/18/13



SUSITNA-WATANA HYDROELECTRIC PROJECT
SUB-ICE CHANNELS CUT THROUGH
INTERFLUVES, SCOTLAND AND
EXAMPLE SUB-ICE CHANNEL MORPHOLOGY

FIGURE
3-1




Source: <http://www.graenslandet.se/en/traces-of-the-ice-age/meltwater-ridges-meltwater-channels-or-glacial-grooves>

Sub-ice channels at Grövelsjön.

DRAFT



Date 10/18/13



SUSITNA-WATANA HYDROELECTRIC PROJECT


EXAMPLE SUB-ICE CHANNELS, GREENLAND

FIGURE

3-2



White arrows denote locations of linear to sub-linear incised creeks that enter at high angles to Seneca Valley and Lake.



Base data from ASTER Global Digital Elevation Model (ASTER GDEM is a product of METI and NASA)

Explanation

Alaska Paleo-Glacier Atlas v. 2 Data
(Kaufman et al., 2011)

Limit of late Wisconsin glaciers


Cosmogenic Exposure Sample Locations

- Dortch et al., 2010a
- Dortch et al., 2010b
- Matmon et al., 2006



Glacial Lake Elevation Extents (meters)

- 800 m
- 975 m

Quaternary fault, solid where well constrained, long dashed line where moderately constrained, short dashed line where inferred (Alaska Division of Geological and Geophysical Surveys, 2012)




79_218900_Alaska_Railbelt/2189_Lineament Report October 2013, modified 12.02.13



Talkeetna fault traces in top panel:

- A** Csejtey et al., 1978
- B** WCC report, 1982
- C** Wilson et al., 2009




SUSITNA-WATANA HYDROELECTRIC PROJECT


WCC TRENCH T-1
LOCATION MAP

FIGURE
4-1

DRAFT




View of WCC T-1 location (marked by tree line), looking slightly east of south.



Very low altitude view of tree line that corresponds to backfilled Trench T-1, with scarp-like feature in mid-background.





View of WCC T-1 looking southwest. Note how the expression of the scarp feature dies out along the projected trend of the feature.




View of WCC T-1 looking northeast.


DRAFT



View looking north-northeast along trend of mapped Talkeetna fault trace with unfaulted volcanic intrusives (Tvu) in the background.



Geology from Wilson et al., 2009



Hillshade from 5-m InSAR data, 2010.

Notes: 1. See Figures A0.2, A0.3, A0.4, and A0.5 for explanation.

DRAFT



SUSITNA-WATANA HYDRO

Clean, reliable energy for the next 100 years.


ALASKA ENERGY AUTHORITY

AEA11-022

16-1401-TM-012014

Appendix A:
Strip Maps and Photographic Documentation
of Lineament Data Presented in FCL (2013)

79_218900_Alaska_Railbelt2189_Lineament Report October 2013, modified 12.16.13
























Explanation	
	Quaternary fault, solid where well constrained, long dash where moderately constrained, short dash where inferred (Koehler et al., 2012)
	Extent of stripmap tile; figure number indicated
	Field work planned in 2013 based on results of TM-8 (FCL, 2013)
	No field work planned in 2013 based on results of TM-8 (FCL, 2013)
	Proposed Watana site

Lineament Groups and Corresponding Figures	
Lineament Group	Appendix A Figure Number
1	A1.1, A1.2
2	A2.1, A2.2
3a	A3a.1, A3a.2
3b	A3b.1, A3b.2
4	None, see TM-8 (FCL, 2013)
5	A5-1.1, A5-2.1, A5-2.2
6	A6.1, A6.2, A6.3, A6.4
7	A7.1, A7.2
8	A8-1.1, A8-2.1, A8-2.2, A8-2.3
9	A9-1.1, A9-2.1, A9-2.2, A9-2.3, A9-2.4
10	None, see TM-8 (FCL, 2013)
11	None, see TM-8 (FCL, 2013)
12a	A12a.1, 12a.2
12b	A12b.1, 12b.2
13	None, see TM-8 (FCL, 2013)
14	None, see TM-8 (FCL, 2013)
15	None, see TM-8 (FCL, 2013)
16	None, see TM-8 (FCL, 2013)
17a	A17a.1, A17a.2
17b	A17b.1, A17b.2, A17b.3
17c	A17c.1, A17c.2
18	None, see TM-8 (FCL, 2013)
19	A19-1.1, A19-1.2, A19-1.3, A19-2.1, A19-2.2, A19-3.1, A19-3.2
20	A20.1, A20.2, A20.3, A20.4, A20.5, A20.6
21a	A21a.1, A21a.2
21b	A21b.1, A21b.2, A21b.3
22	A22.1, A22.2
23	A23.1
24	None, see TM-8 (FCL, 2013)
25	None, see TM-8 (FCL, 2013)
26	A26.1, A26.2
27	A27-1.1, A27-2.1, A27-3.1, A27-3.2
Broad Pass area	Plate A-BP, A-BP.1, A-BP.2, A-BP.3
Castle Mtn. fault extension	Plate A-CME, A-CME.1, A-CME.2
Clearwater Mtns. area	Plate A-CWM, A-CWM.1, A-CWM.2, A-CWM.3

79_218900_Alaska_Railbel/2189_Lineament Report October 2013, modified 01.06.14


























This explanation applies to all figures and plates in Appendix A.

Geologic Units from OFR 09-1108 (Wilson et al., 2009)













	Water, ice field, or glacier
Unconsolidated Deposits	
	Surficial deposits, undivided
	Alluvium along major rivers and in terraces
	Landslide and colluvial deposits
	Glacial deposits, undivided
	Young moraine deposits
	Major moraine and kame deposits
	Glacioalluvium
	Outwash in plains, valley train, and fans
	Glacioestuarine deposits
Sedimentary Rocks	
	Sedimentary rocks, undivided
	Kenai Group, undivided
	Tsadaka Formation
	Chickaloon formation
	Matanuska formation
	Turbiditic sedimentary rocks of the Kahiltna flysch sequence
	Undivided Chinitna and Tuxedni formations
	Naknek Formation, undivided
	Talkeetna Formation, undivided
	Limestone and Marble
	Eagle Creek Formation, marine argillite and limestone

Note: For full explanation of geologic units see USGS OFR 09-1108 and USGS OFR 98-133.



Igneous Rocks

Volcanic and Hypabyssal Rocks	
	Tertiary volcanic rocks, undivided
	Felsic volcanic and sub-volcanic rocks
	Mafic volcanic rocks
	Dikes and sills
	Nikolai Greenstone and related rocks
	Slana Spur Formation, volcanoclastic rocks
	Station Creek Formation andesitic volcanic rocks
Plutonic Rocks	
	Intrusive rocks, undivided
	Granitic rocks
	Granitic rocks of Paleocene age
	Biotite-hornblende-granodiorite
	Granitic rocks, undivided
	Granodioritic rocks
	Granodiorite
	Trondhjemite
	Diorite, gabbro, picrite, and pyroxenite sill and dike swarm complex
	Quartz diorite, tonalite, and diorite
	Granodiorite and quartz monzonite
Melange and Metamorphic Rocks	
	Gneiss
	Plutonic and metamorphic rocks, undifferentiated
	Amphibolite
	Marble
	Metabasalt and slate
	Basaltic to andesitic metavolcanic rocks
	Metamorphosed Skolai Group


Geologic Units from OFR 98-133 (Wilson et al., 1998)

	Ice fields or glaciers
	Water
	Surficial deposits, undifferentiated
	Tertiary volcanic rocks, undivided
	Hypoabyssal felsic and intermediate intrusions
	Granitic and volcanic rocks, undivided
	Granite and granodiorite
	Phyllite, pelitic schist, calc-schist, and amphibolite of the MacLaren metamorphic belt
	Granitic rocks
	Kahiltna flysch sequence
	Calcareous sedimentary rocks
	Metavolcanic and associated metasedimentary rocks













Tectonic Features from WCC report (WCC, 1982)

-  Detailed feature, from site-specific maps
-  Regional feature, from small-scale maps




For completeness, features from both regional and detailed scale figures have been included. The location of regional features may not always be accurate and the detailed features may be limited to the extent shown on original figures.

 Location of trench T-2 (shown on Figures A14 and A16)




Faults Compiled by FCL (Wilson et al., 1998; Wilson et al., 2009; Williams and Galloway, 1986; Clautice, 1990; Clautice, 2001; Csejtey, 1978; Kachadoorian, 1979; Smith, 1988)

-  Fault, approximate
-  Fault, inferred or queried
-  Fault, certain
-  Fault, concealed
-  High-angle reverse fault, approximate
-  High-angle reverse fault, certain
-  High-angle reverse fault, concealed
-  High-angle reverse fault, inferred or queried
-  Thrust fault, approximate
-  Thrust fault, certain
-  Thrust fault, concealed
-  Lineament

Hydrographic Features from National Hydrography Dataset, 2000, 1:24,000 scale

-  Stream
-  Ice mass
-  Lake or pond

Other Items

-  Location of photograph taken during 2013 field reconnaissance, labeled with photo ID and showing view direction
-  GPS waypoint
-  GPS track line, July 2013

79_218900_Alaska_Railbelt/2189_Lineament Report October 2013, modified 01.06.14

Attributes of lineaments mapped by FCL (2013) that apply to all figures and plates in Appendix A

Reconnaissance (INSAR)

- 1 - 5
- 10
- 77
- 88

Detail (LiDAR)

- 1 - 5
- 10
- 77
- 88

Lineament Groups

- 17a Lineament group mapped for this study coinciding with previously mapped fault or lineament
- 25 No previously mapped fault or lineament coincides with lineament group

Explanation for relevant geologic units of Williams and Galloway (1986) shown on Figure A20.5 and A23.1

Geologic Units

Bottom deposits of 914 - 975 m lake
Overprint denoting glacial drift that is mantled by bottom sediments of glacial lake that extended to 914 - 975 m above modern sea level, largely confined to middle Susitna valley, above ice dam below Fog Lake (off map) and apparently bounded on east and south side by glacier ice. Does not cover late(st) Wisconsin (last major) morainal systems. No shoreline features are mapped.

Bottom deposits intermediate (777 - 747) lake
Overprint denoting bottom deposits of a local lake that covered melting glacier ice between Tyone Lake and Lake Louise, apparently behind Tyone Spillway, and drained as the elevation of the spillway was cut down from 777 m to 747 m above sea level while stagnant ice was still in valley bottom.

Bottom deposits of last regional lake
Overprint denoting drape of bottom deposits over drift and thick lake sediments that persisted in Copper River drainage basin from just before deposition of Old Man moraines to a time when glaciers had retreated to within 16 to 24 km of present glaciers: older than 13,000 years.

Symbols

- AA Location and letter designation of radiocarbon-dated stratigraphic section in accompanying text.
- Ice boundary, morainal ridge, kame terrace, delta, or other ice contact feature marking edge of glacier: hachures toward glacier.
- Shoreline of regional lake: mapped for the lake in Copper River basin where at 747 m (maximum elevation); the elevation to which Tyone Spillway was eroded, and successively lower levels in the northern part of area between 747 m and 701 m above sea level. Lesser recessional shorelines mapped by Nichols and Yehle (1969) not shown.
- Upper limit of post-glacial (Holocene, in part) shoreline of Tazlina Lake from elevation 564 m down to present lake level 544 m caused by lowering of lake as Tazlina River has deepened its canyon.
- Delta of glacial lake, including those of modern glacial lakes such as Tazlina Lake.
- Linear or drumlinoid feature, due to ice scour, direction of ice movement indicated by arrow.
- Spillway for glacial meltwater, including that stored in large glacial lakes.
- Contact between map units where not glacial boundary, most commonly between different levels of lake deposits.
- Active (?) fault, lower Sonoma Creek, offsetting unconsolidated deposits.
- Location of selected erratic boulders, mountain top erratic stones transported by glaciers, e.g. Sheep Mountain; many occurrences on mountains lower than 1829 m not shown.

Attribute	Cross Section Morphology*	Description	Examples
1		Linear break-in-slope bisecting a planar surface	Uphill- or downhill-facing scarps, ateral moraines or kame deposits along lateral margins of valley glaciers
2		Abrupt changes in slope adjacent to otherwise relatively horizontal (and planar) surfaces	Linear range fronts, faceted ridges, terrace risers, steep downstream faces of roushe mountonees
3		Linear U-shaped trough	Glacial valleys, ice-scoured flutes, flood-scoured flutes,
4		Linear V-shaped trough	Active stream channels
5		Linear ridges	Drumlins, water-scoured terrain, eskers
6 (also 77)	n/a	A series of aligned features	Could include attributes #1 -5 above and/or aligned saddles, tonal lineaments, etc.
66	n/a	Data artifacts	Linear seams between data sets collected on different dates
88	n/a	A series of aligned features, which are too small to individually map at the given scale	Could include features with attributes #1-5 above and/or aligned saddles, tonal lineaments, etc.
99	n/a	A line which encloses a broad expanse of features all having the same orientation	An area of jointing or of glacial striae all having the same, parallel orientation
10	n/a	Anthropogenic lineaments	Roads, rail roads, power lines and other linear clearings, etc.

Notes: *Arrow points to location of the mapped feature.

79_218900_Alaska_Railbelt2189_Lineament Report October 2013, modified 10.18.13

Explanation for relevant geologic units of Smith et al. (1988) shown on Figure A21b.1

UNCONSOLIDATED DEPOSITS

Alluvial deposits

Qa FLOODPLAIN ALLUVIUM - Unconsolidated deposits in modern stream drainages. Material ranges from coarse, unsorted gravel in highland valleys to finely bedded silt in large river drainages.

Glacial deposits

Qdt₃ TILL OF LATE WISCONSIN AGE - 11,800 to 25,000 yr B.P.


Qdt₂ TILL OF EARLY WISCONSIN AGE - 40,000 to 75,000 yr B.P.


Ks SCHIST - Medium- to coarse-grained biotite-plagioclase-quartz schist with local garnet and feldspar porphyroblasts to 0.5 mm. Dominantly gray or brown weathering. Includes local horizons that contain randomly oriented hornblende on foliation surfaces. Stippled pattern near intrusive contacts indicates hornfelsed zone in schist. K-Ar age of 57.2 m.y. was obtained from biotite in this unit in the adjacent Healy A-1 Quadrangle (Smith, 1981).


Kp PHYLLITE - Silver-gray, biotite-bearing phyllite with biotite porphyroblasts to 2mm long; locally calcareous. Minor compositional banding with more quartzose layers parallel to foliation. Biotite yielded K-Ar age of 53 ± 1.6 m.y. (loc. 3 on map; Turner and Smith, 1974). Grades into amphibole-bearing phyllite (Khp) unit.


Kph AMPHIBOLE-BEARING PHYLLITE - Medium to dark gray spotted phyllite with planar laminations. Spotted with porphyroblastic biotite. Interlayered with beds that contain randomly oriented amphibole on foliation surfaces. Amphibole prisms commonly 0.5 to 3 mm long. K-Ar age of actinolitic hornblende from this unit in Healy A-I Quadrangle is 64.1 m.y. (Smith, 1981).

MAP SYMBOLS

 Contact - dashed where approximately located ; dotted where concealed; queried where inferred

 High-angle fault - dashed where approximately located; dotted where concealed; queried where inferred. D, downthrown side; U, upthrown side


 Thrust fault - dashed where approximately located. Sawteeth on upper plate. Arrow indicates dip of fault


 Lineament - inferred from aerial photographs, may represent fault


Modified from selected portion of Smith et al. (1988) explanation


Explanation for relevant geologic units of Reger (1990) shown on Figure A21a.2

GLACIAL LIMITS


 Glaciation of unassigned age, dashed where discontinuously mapped


 Glaciation of Illinoian age, dashed where discontinuously mapped

 Glaciation of late Wisconsin age, dashed where discontinuously mapped


 Glaciation of Holocene age, dashed where discontinuously mapped


OTHER FEATURES


 Prominent meltwater drainage channel

 Radiocarbon sample locality


PROMINENT WAVE-CUT SCARPS


 3,700-ft (1,120-m) lake, dashed where discontinuously mapped, dots on descending scarp


 3,650-ft (1,110-m) lake, dashed where discontinuously mapped, open triangles point down descending scarp

 3,400-ft (1,030-m) lake, dashed where discontinuously mapped, solid triangles point down descending scarp


AREAS INUNDATED BY GLACIER-DAMMED LAKES


 3,700-ft (1,120-m) lake


 3,650-ft (1,110-m) lake

 3,400-ft (1,030-m) lake

Explanation for relevant geologic units and features from Acres, 1982 shown on Figure A6.1

 Contact

 Thrust fault

 Shear

QUATERNARY

Qa Alluvium, alluvial terraces and fans

Qid Ice disintegration deposits

Qt Till

Qo Outwash

TERTIARY

Tsu Conglomerate, sandstone and claystone

MESOZOIC



TRIASSIC

TRvs Basaltic metavolcanic rocks, metabasalt and slate



DRAFT

Explanation

Lineaments, Faults, Contacts, Synclines, and Anticlines




DRAFT




Notes: 1. See Figures A0.2, A0.3, A0.4, and A0.5 for explanation.
 2. Geology by Wilson et al., 1998

DRAFT


79_218900_Alaska_Railbelt/2189_Lineament Report October 2013, modified 10.18.13



View looking northeast from location A towards the confluence of the Jack River and the East Fork Jack River. Arrows point along the alignment of mapped lineaments. Note absence of linear expression in Quaternary deposits.





View looking southwest from location B along alignment of linear features. Arrows indicate the alignment of the mapped lineaments.




View looking southwest from location C at a detailed view of aligned uphill-facing scarps. Note Thf contact is up-slope from the scarp in the distance.

DRAFT



Notes: 1. See Figures A0.2, A0.3, A0.4, and A0.5 for explanation.
2. Geology by Wilson et al., 1998.




SUSITNA-WATANA HYDROELECTRIC PROJECT
LINEAMENT GROUP 2
MAP DATA

FIGURE
A2.1



DRAFT




Photograph taken from location A looking east-northeast. Arrows show the alignment of FCL-mapped lineament. Note lack of apparent deformation in bedrock exposure along Jack River.



DRAFT



Notes: 1. See Figures A0.2, A0.3, A0.4, and A0.5 for explanation.
2. Geology by Wilson et al., 1998.




SUSITNA-WATANA HYDROELECTRIC PROJECT
LINEAMENT GROUP 3a
MAP DATA

DRAFT


FIGURE
A3a.1



View looking east at likely solifluction-related scarps on hillside that correspond with mapped lineaments. Large arrows point along lineaments.





View looking west along 3a lineament expressed as sharp ridge within Kahlitna flysch (KJf). Apparent color change and topographic expression may suggest a geologic structure, however, none were previously mapped. The feature may be a result of weathering because of lithologic change within the flysch.



View looking east past ridge, with unfaulted Quaternary sediments in the foreground and far distances.

DRAFT



Notes: 1. See Figures A0.2, A0.3, A0.4, and A0.5 for explanation.
2. Geology by Wilson et al., 1998.



SUSITNA-WATANA HYDROELECTRIC PROJECT
LINEAMENT GROUP 3b
MAP DATA

DRAFT

FIGURE
A3b.1




View looking west along north-facing escarpment in Eocene granitics.



View looking east along lower talus scree field that shows decreasing relief at west end of lineament 3b.





View looking west along lineament 3b projection. South-facing escarpment indicates a reversal in kinematic morphology.





View looking west along lineament 3b projection. Holocene rock glaciers are not offset, and lineament is expressed as a linear valley.


DRAFT



Notes: 1. See Figures A0.2, A0.3, A0.4, and A0.5 for explanation.
2. Geology by Wilson et al., 2009.




Notes: 1. See Figures A0.2, A0.3, A0.4, and A0.5 for explanation.
2. Geology by Wilson et al., 2009.




SUSITNA-WATANA HYDROELECTRIC PROJECT
LINEAMENT GROUP 5
MAP DATA

DRAFT


FIGURE
A5-2.1



View looking west at eastern part of apparent side hill bench.





View looking west along ice-scoured terrain, with the Indian River flowing from right to left.




View of linear gullies developed on bedrock slope. Mapped lineament approximately shown.




View of drainage with mapped lineament approximately shown.



Notes: 1. See Figures A0.2, A0.3, A0.4, and A0.5 for explanation.
 2. Data frame has been rotated 45° east of north.
 3. Geology from Acres, 1982 (top) and by Wilson et al., 2009 (bottom)




View looking west along oblique to projection of Talkeetna fault



View looking east along lower river bank at apparent alternation zone distinguished by color contrast, possible juxtaposition of Triassic metabasalts and undifferentiated Tertiary sediments. This location is east of the mapped projections of the Talkeetna fault.




View looking east at apparent flat-lying contact between Quaternary lake sediments (above) and Quaternary till (below). Arrows point to contact.



View looking west at projected trace of Talkeetna fault whose ground expression is absent in Quaternary surface.


DRAFT



View looking south at erosion-resistant ridge of Tertiary sediments whose beds dip gently to the northwest but appear undisrupted.



View looking west at apparently northwest-dipping beds in Tertiary sediments, relatively consistent with northwest dips measured by WCC (1982) in Tertiary sediments along west bank Watana Creek.



View looking north-northeast past ridge, with flat and apparently undisturbed Quaternary sediments in the background.



View looking west at bedded (lake?) stratigraphy exposed in eroding bluff. Beds appear relatively horizontal, but may have a sense of non-planar geometry because of semi-circular outcrop. Note fallen trees that indicate erosion/slope movement.


DRAFT

I)





View looking north at linear esker nearly coincident with map projection of Talkeetna fault.

J)




View looking at shallow soil pit dug in esker crest. Upper black, gray, and reddish soil layers are Holocene tephras. Scale is in centimeters; the upper 45 centimeters of the pit are in view.

DRAFT



Notes: 1. See Figures A0.2, A0.3, A0.4, and A0.5 for explanation.
 2. Data frame has been rotated 45° east of north.
 3. Geologic map by Kline et al., 1990.




SUSITNA-WATANA HYDROELECTRIC PROJECT
 LINEAMENT GROUP 7
 MAP DATA

DRAFT
 FIGURE
 A7.1



View looking at color contrast at previously mapped bedrock fault.



View looking west down-valley at apparent undeformed glacial sediments.



View looking up-valley at incised drainage that coincides with mapped lineament and previously mapped fault.



View looking down-valley from the top of the drainage seen in Photograph C.